

THE INDIVIDUALITY OF THE GERM-NUCLEI
DURING THE CLEAVAGE OF THE EGG OF
CRYPTOBRANCHUS ALLEGHENIENSIS.

BERTRAM G. SMITH.

Early observers of the process of fertilization described the meeting of the sperm-nucleus and the egg-nucleus, and their complete fusion to form a single zygotic nucleus, the primary nucleus of the embryo. Later it was found that in many cases, though apparently not in all, the two germ-nuclei merely become apposed without fusion. In tracing their further history it was shown, in certain cases, that the germ-nuclei give rise to two independent groups of chromosomes which separately enter the equatorial plate and whose descendants pass separately into the daughter-nuclei. "Later observations have given the strongest reasons for believing that, as far as the chromatin is concerned, a true fusion of the nuclei never takes place during fertilization, and that paternal and maternal chromatin *may* remain separate and distinct in the later stages of development—possibly throughout life" (Wilson, 1900, p. 204). At the present time there is much to warrant the belief that a fusion of maternal and paternal chromatin never takes place in the somatic cells, while in the lineage of germ cells nothing approaching fusion occurs until a pairing of maternal and paternal chromosomes, called synapsis, takes place in preparation for the maturation divisions; then for the first time maternal and paternal chromosomes are brought together in intimate and orderly union, in some cases amounting to fusion.

This conclusion, which is of the most fundamental importance since it vitally concerns the mechanism of heredity, is based largely on indirect evidence; for in almost every case apparently insuperable obstacles have been encountered in the attempt to trace the respective maternal and paternal chromatin-complexes through the resting stage of the nuclear cycle. In only a very

few cases has the individuality of the germ-nuclei during this stage been demonstrated with any considerable success; the observations of Häcker (1892), Rückert (1895) and Conklin (1901) are of most importance in this connection and will be discussed later.

In the developing egg of the amphibian *Cryptobranchus allegheniensis* I have found material very favorable for the study of this problem. The early cleavage nuclei are invariably double throughout the entire resting stage, each consisting of two separate though closely apposed nuclear vesicles, and the origin of each vesicle from a single germ-nucleus has been clearly traced. Indeed, my observations indicate that typically the germ-nuclei maintain their individuality throughout cleavage, though certain irregularities, real or apparent, occur with increasing frequency in the later segmentation stages.

TECHNIQUE.

Most of the sections used in making the observations recorded in the present paper were cut from eggs killed with bichromate-acetic-formalin. This mixture, made up in the right proportions (Smith, 1912, I.), preserves the form of the egg perfectly in all stages, enables it to be sectioned by the paraffin method, and gives very satisfactory histological and cytological results. For comparison, sections were made of a few eggs killed in Zenker's fluid, also in corrosive-acetic-formalin. These mixtures, particularly Zenker's fluid, gave excellent results in the fertilization and early cleavage stages, but in the blastula and gastrula stages their use is not advisable since the roof of the blastocœle or gastrocœle collapses during the process of infiltration with paraffin.

Most of the observations were made on sections of eggs stained *in toto* with Grenacher's borax-carmine, differentiated in acid alcohol, and counterstained on the slide with a mixture of Lyons blue and picric acid. This treatment leaves the chromatin stained red, the yolk granules green, the centrospheres and cytoplasmic structures blue. The method has the advantage of rapidity of manipulation, an important consideration since a

large number of slides are required. For the main purposes of this investigation the differentiation obtained by this method, used with care, is entirely adequate, but as a precaution against possible inaccuracy a parallel series was stained with Heidenhain's iron hæmatoxylin, which as might be expected gave better preparations, and is especially useful for the study of chromosomal vesicles.

All the figures excepting those of Plate IX. were outlined by means of a camera lucida and drawn with a magnification of about 600 diameters; in the process of reproduction they were reduced one third, and thus appear with a magnification of about 400 diameters. In all drawings of horizontal sections the orientation with reference to the plane of first cleavage is the same, and conforms to the diagrams in Plate IX.

In most cases the double nature of the cleavage nuclei appears to best advantage in horizontal sections, and might readily be overlooked were observations confined to vertical sections. It seems likely that most investigators studying the amphibian egg have confined their observations to vertical sections, since these give a better idea of the gross structure; and this circumstance leads us to suspect that conditions such as are here described for *Cryptobranchus* may occur in other amphibians and need only to be revealed by the study of horizontal sections.

OBSERVATIONS.

The early history of the fertilization stage of *Cryptobranchus allegheniensis* has been given in a previous paper (Smith, 1912, I.). The superficial cleavage pattern has been described in another contribution (Smith, 1912, II.).

At the time of the meeting of the germ-nuclei it is usually impossible to tell with certainty which is the egg-nucleus and which is the sperm-nucleus, but this circumstance does not affect the validity of the results recorded in this paper. The germ-nuclei do not fuse, but come to lie side by side with nuclear membranes intact (Figs. 1-4). During the long resting stage that precedes the formation of the first cleavage spindle, each germ-nucleus maintains strict individuality: there is close asso-

ciation, but no actual union, and certainly no mingling of structural contents. Hence, although it is convenient to employ such terms as "union of the germ-nuclei," and to speak of the "cleavage-nucleus" or "segmentation-nucleus" as if it represented a single entity, it should be understood that such expressions are loosely used, and the word "fusion" in this particular case is altogether inapplicable. It is the purpose of this paper to show that this unity or individuality of the respective germ-nuclei is not confined to the fertilization stage, but persists with great regularity during early cleavage, and with some apparent irregularities throughout the later cleavage and even into the gastrula stage.

The length of time intervening between the meeting of the germ-nuclei and the rupture of their nuclear membranes during the formation of the first cleavage spindle must be considerable, for resting germ-nuclei have been observed in eggs varying in age from twelve to twenty-three hours after fertilization. The precise time of formation of the first cleavage spindle has been determined in only one egg, which was killed twenty-six hours after artificial fertilization.

Since the germ-nuclei usually come to lie in the same horizontal plane, the most fruitful observations are obtained through the study of horizontal sections. The two asters, formed by the division of the single aster accompanying the sperm-nucleus on its journey to meet the egg-nucleus, take up positions on opposite sides of the resting germ-nuclei, in the same horizontal plane, and close to the region of contact of the germ-nuclei (Figs. 3 and 4). The initiation of the process of mitosis as evidenced by the appearance of distinct chromosomes and the rupture of the nuclear membranes does not take place in the two germ-nuclei simultaneously; one of the germ-nuclei becomes active somewhat in advance of the other (Figs. 5 and 6), thus furnishing evidence of some degree of physiological as well as structural independence.

In the fully-formed mitotic figure the two groups of chromosomes, of maternal and paternal origin respectively, remain visibly distinct (Fig. 7). This has been determined beyond a

doubt in a number of cases, with no conflicting evidence. In the anaphase each of the original chromosome groups has divided equally (through the splitting of individual chromosomes), the two halves going to different poles of the spindle. Each daughter-nucleus thus receives two distinct groups of chromosomes, derived from the egg-nucleus and the sperm-nucleus respectively. Although in the anaphase and early telophase (Fig. 8) the paired groups of chromosomes are not always very sharply separated, yet there is no mingling of maternal and paternal chromosomes. Finally, in the newly-formed daughter-nucleus each parental group of chromosomes becomes metamorphosed into a separate nuclear vesicle (Figs. 9 and 10). Except for a diminution in size, which is afterwards partially made good by growth, each daughter-nucleus closely simulates the original cleavage-nucleus with its two resting germ-nuclei.

The precise stage with newly-formed vesicular daughter-nuclei has not been found in horizontal sections, hence all the figures of this stage are taken from vertical sections (Figs. 9-12). During the late phases of mitosis the two daughter-nuclei move some distance apart; their path of separation is marked by a broad trail of cytoplasm sharply defined from the surrounding yolk. The division of the centrosphere or "attraction sphere" takes place shortly before the daughter-nucleus assumes the vesicular condition. The first cleavage furrow is not yet formed, but its position, in a plane at right angles to the path of separation of the two daughter-nuclei and midway between them, is visibly indicated by a condensation of cytoplasm in the blastodisc.

Aside from the fact that the two nuclear vesicles of a newly-formed daughter-nucleus are always situated approximately equidistant from the plane of first cleavage, the orientation of these nuclear vesicles with reference to each other is decidedly variable. Since the two original germ-nuclei at the time of meeting usually lie in the same horizontal plane (Figs. 2-4), we should expect the same thing of the two vesicles of a daughter-nucleus; but in the preparations studied these were more often found to lie in a plane oblique to the horizontal, and sometimes one vesicle lies directly above the other (Figs. 9 and 10). It should be remembered,

however, that the original germ-nuclei do not always lie in the same horizontal plane (Fig. 1). Moreover, the stage with newly-formed daughter-nuclei was not represented in horizontal sections, which are naturally most favorable for the discovery of nuclei consisting of vesicles lying in the same horizontal plane. There is no external landmark by means of which this stage may be recognized, and it is passed through very rapidly; in the making of horizontal sections the available material was exhausted without finding the precise stage desired.

In the study of the subsequent behavior of the nuclei the early cleavage furrows serve as convenient landmarks for orientation, and it is necessary to keep in mind the elementary fact that the second cleavage furrow forms at right angles to the first, and that a similar alternation in the direction of successive cleavage furrows is characteristic of the entire process of cleavage. Of more immediate importance to us is the correlated fact that the second nuclear division occurs in a direction at right angles to the first nuclear division, and that this alternation in the direction of successive nuclear divisions is repeated throughout cleavage. To be sure, this is strictly true only in the early cleavage stages, for there is increasing irregularity in the direction of both cleavage furrows and nuclear divisions throughout the later stages of cleavage. Incidentally, it should be noted that nuclear division always precedes the formation of the corresponding cleavage furrow.

In preparation for the second nuclear division, the two asters of a single daughter-nucleus take up positions on opposite sides of the nucleus, in the same horizontal plane and with a line connecting them parallel to the first cleavage furrow which is now usually in process of formation (Figs. 12-20; 23-26). In case the two nuclear vesicles of a single daughter-nucleus are not already ranged midway between the two asters, they rotate to this position (compare Figs. 15-16 with Figs. 17-18). Typically, the two nuclear vesicles come to lie in the same horizontal plane (Figs. 17-20, 23-26); but this is not the invariable position, for in three different eggs they were found to lie one above the other, as exemplified in Figs. 21 and 22. In any case the final

position assumed insures an equal division of maternal and paternal chromatin-groups in the ensuing mitosis. Thus the second cleavage mitosis (Figs. 27 and 28) is in all essential respects a repetition of the first, though the direction of the spindle is different. Throughout the entire segmentation period there is nothing in the behavior of the nuclei more impressive than the precision and regularity with which the two nuclear vesicles of a single nucleus are ranged side by side in the plane of the equatorial plate in preparation for the formation of the spindle.

In the resting stage following the second mitosis the two nuclear vesicles of each nucleus usually lie in the same horizontal plane (Figs. 29, 30, 32); their entire orientation indicates their derivation from the two groups of chromosomes at one pole of the spindle in the anaphase and early telophase of the preceding mitosis (Fig. 28). Occasionally, though not so often as in the preceding resting stage, one nuclear vesicle is found situated above the other (Fig. 31).

In *Cryptobranchus* the third cleavage furrows are vertical and intersect the second furrows at some distance from the first (Plate IX., Fig. 45, *J*). Thus the third cleavage furrows are nearly parallel to the first, but diverge slightly from it.

In preparation for the third cleavage mitosis the two asters take up positions on opposite sides of the nucleus, in such a situation that a line connecting them is nearly parallel to the plane of second cleavage (Figs. 33, 34; Plate IX., Fig. 45, *H*). In all cases the two nuclear vesicles entering the mitotic figure are ranged equidistant from the two asters (Figs. 33, 34, 35), and in a slightly earlier stage may be found rotating to this position (Fig. 32). Thus in the third cleavage mitosis (Fig. 36) the segregation and equal division of maternal and paternal chromosome groups is maintained. The direction of the axis of the spindle is at right angles to the preceding mitosis, and parallel to the first (Plate IX., Fig. 45, *H* and *I*). Upon the completion of mitosis the third cleavage furrows form as described in the preceding paragraph. At this time the four nuclei on the same side of the second cleavage furrow sometimes lie precisely in

the same vertical plane, so that they are all visible in a single section taken parallel to the second cleavage furrow.

The fourth cleavage furrows combine to form a circular, or more often oval, figure with its longer axis coinciding with the plane of second cleavage. This circular cleavage divides the egg very unequally, cutting off eight micromeres from eight comparatively large macromeres. Preparation for this cleavage involves rotation of the two nuclear vesicles of a single daughter-nucleus of the third cell division from their original position as shown in Fig. 37 to the position shown in Fig. 38. This latter orientation insures the maintenance of the segregation of the germ-nuclei in the fourth mitosis (Figs. 39-42). As in the previous divisions, the maternal and paternal chromosome groups have here been traced through the various phases of mitosis to the late anaphase and early telophase, where they are still distinctly separate. The position of the vesicles of the newly-formed daughter-nucleus (Figs. 43 and 44) indicates that one is of maternal origin, the other paternal. The results as regards the orientation of the nuclei up to this point are summarized in the diagrams of Plate IX., Fig. 45.

In the later cleavage divisions there is increasing irregularity in the direction of spindle axes and cleavage furrows, but this does not constitute any serious obstacle to tracing the history of the germ-nuclei since the really important point is the orientation of the nuclear vesicles with respect to the axis of the spindle. As already indicated, this is invariably such as to insure the segregation and equal representation of maternal and paternal chromosome groups in the daughter-nuclei. As the nuclei become smaller, it sometimes is difficult to distinguish the two groups of chromosomes in the late phases of mitosis, but comparison with slightly earlier and later phases leaves no doubt as to the continuity of the respective germ-nuclei. Until an advanced cleavage stage, the double structure of the resting nuclei is almost always clearly demonstrable. As the nuclei become more numerous, it becomes easier to find nuclei cut in such a manner as to show distinctly the double structure. In the more active regions of the egg, the synchronism of division

of the different cells is maintained to the late blastula stage, so that there is little likelihood of confusing cell-generations. Between micromeres and macromeres there is a fine gradation in phases of the nuclear cycle, so that a close series may be readily obtained. The individuality of the germ-nuclei has been traced without a break to an advanced cleavage stage, and it would not be difficult to illustrate this entire result with an adequate series of figures, but such a procedure would require an enormous amount of time and unduly extend the limits of this paper.

Beginning with about the seventh cell generation certain irregularities, real or apparent, occur with increasing frequency to mar the mechanical precision of the events thus far described in the history of the germ-nuclei. In the resting stage the two nuclear vesicles of a single nucleus are often deeply lobed, sometimes in such a manner as to give the impression that the nucleus is made up of many vesicles. In the late telophases the nucleus is often really made up of several or many separate though closely aggregated vesicles, but this is undoubtedly a part of the usual process of mitosis, as a more searching examination of these late phases throughout cleavage will readily show.

The transformation of the daughter-nucleus into the vesicular condition must take place with considerable rapidity, especially in the early cleavage stages, for some phases of this process have not been observed earlier than the fourth mitosis, and only occasionally in the fourth and fifth mitoses. In more advanced cleavage there is little difficulty in finding material representing the entire history of the telophase.

In the early anaphase, the long thread-like chromosomes are V-shaped, with the limbs of the V almost parallel and very straight; the apex is directed away from the equatorial plate. In a slightly later phase, which we may call the late anaphase, the chromosomes are still roughly V-shaped but finely undulating except at their free ends. A little later (early telophase) each chromosome becomes much convoluted, often coiled in a loose irregular spiral; this change begins at the apex and proceeds toward the free ends. The next step consists of the metamorphosis of each chromosome into a chromosomal vesicle, a

process which I have been able to follow in considerable detail, but will not attempt to describe here. About this time the centrosphere or "attraction sphere" divides.

During the anaphase and early telophase the maternal and paternal chromosome groups are almost always clearly separated, sometimes with a broad space between the two groups. The segregation of the chromosomal vesicles into two groups, maternal and paternal respectively, is usually evident. In the late telophase the chromosomal vesicles of each group unite to form separate nuclear vesicles which for a time show indications of their manner of origin through the persistence of lobes separated by deep clefts. The inner boundaries of the chromosomal vesicles persist for a time as partitions in the nuclear vesicles, but finally disappear wholly or in part. In the late blastula the vesicular nucleus in many cases retains a lobed structure throughout the resting stage. It is possible that in some cases the fusion of chromosomal vesicles to form vesicular nuclei of the usual duplex type is never completed, for in the late blastula and early gastrula nuclei consisting of three, four or even more well-rounded vesicles are not uncommon.

How much longer and to what extent during the development of the embryo a strict separation of the germ-nuclei is maintained I have not attempted to determine, but it is not necessary to conclude that it continues throughout the life of the organism, either in the somatic or the germ cells. In the discussion it will be shown that the individuality of the germ-nuclei is not incompatible with a mingling of maternal and paternal chromosomes, and there is no biological necessity for a strict separation of the germ-nuclei in any stage from fertilization to synapsis.

DISCUSSION.

In 1875 Oscar Hertwig and Hermann Fol showed that the fertilized egg contains two nuclei, one belonging to the egg itself and the other introduced by the spermatozoön. While the earliest observers of the process of fertilization, notably Auerbach, Strasburger and Hertwig, described the complete fusion of these germ-nuclei to form the first embryonic nucleus, called by

Hertwig the cleavage-nucleus or segmentation-nucleus, later observations showed that such is not always the case. In 1881 Mark demonstrated that in the slug *Limax* the two germ-nuclei do not fuse; after coming together they persist during the formation of the two cleavage centers, then their membranes gradually disappear. Two years later Van Beneden showed that in *Ascaris*, not only do the germ-nuclei become apposed without fusion, but each gives rise to an independent group of chromosomes which contribute equally to the formation of the daughter-nuclei. Thus the foundation of the doctrine of the biparental character of the nuclei of sexually-produced organisms was laid down by Van Beneden. In many animals, and in some plants (gymnosperms), the independence of the formation of the maternal and the paternal chromosome groups following fertilization has been established by direct observation, though the demonstration has seldom been carried beyond the first cleavage stage. On the other hand there are some animals, and many plants, in which the germ-nuclei meet and fuse while in the resting condition, so that in the chromatin of the resulting nucleus maternal and paternal contributions cannot be readily distinguished.

Häcker (1892 and 1895) and Rückert (1895) found that the germ-nuclei of *Cyclops* do not fuse but preserve their individuality throughout at least a considerable period of the cleavage of the egg. In mitosis the two groups of chromosomes, of maternal and paternal origin respectively, remain distinct and bilaterally distributed, while each resultant daughter-nucleus in the resting stage consists of two closely apposed but structurally separate vesicles.

In 1901 Conklin described the double structure of the cleavage nuclei of *Crepidula* in certain stages of the nuclear cycle, and pointed out that the two halves which at times appear as distinct entities are almost certainly to be regarded as of maternal and paternal origin respectively. "This separateness is most easily observed in the telophase of each division, though in some cleavage cells it may be seen in the prophase also, or even throughout the resting period. At the time when the daughter-nuclei are being formed the chromosomal vesicles fuse into two groups

which are closely pressed together but still separated by a partition wall, as Rückert has shown to be the case in *Cyclops*. Gradually this partition wall disappears, being preserved longest on that side of the nucleus nearest the centrosome. Here a groove is formed on one side of the nucleus which marks the line of contact between the two halves. In some cleavage cells this groove is visible throughout most of the resting period; in others it disappears during the greater part of the resting period, though it may reappear in the following prophase; in all cases, however, the partition wall and groove reappear in the next succeeding telophase, when it is again formed in the manner described above. I have observed the double character of the nucleus in the telophase of every cleavage up to the 29-cell stage, and in several of the later cleavages up to the 60-cell stage, though it becomes more difficult to see as the nuclei grow smaller. . . . It still remains to show that these double nuclei really represent the egg and sperm nuclei which have not yet lost their individuality. This cannot be demonstrated in *Crepidula*, for the reason that this double character is not apparent at every stage in the nuclear cycle, but it is extremely probable" (Conklin, 1901). Additional observations recorded by Conklin in support of his interpretation may be summarized as follows: (1) In the metaphase of the first cleavage division the maternal and paternal germ-nuclei are represented by separate groups of chromosomes; in the early anaphase these groups of chromosomes can no longer be distinguished, but the nuclei are clearly double in the immediately following late anaphase and telophase, and the position of the partition wall in these double nuclei corresponds to the plane of contact between the germ-nuclei. (2) The groove which is found on one side of the nucleus in the telophase of the first cleavage mitosis persists well into the resting stage, and a corresponding groove is found in the same position in the prophase of the second mitosis. The central spindle for the second cleavage mitosis lies in this groove, and thus the amphiaster actually lies in the only plane in which it would be possible to halve the two parts of the double nuclei. Although cleavage divisions successively alternate in direction, unequal division of the double nuclei is

prevented either by a rotation of the nucleus during the resting stage, or by a rotation of the spindle in the early stage of mitosis.

(3) In certain abnormal cleavages the double nuclei are really two entirely separate nuclei lying side by side within a single cell.

(4) In each of the germ-nuclei, before they come into contact, there is a single nucleolus; these nucleoli disappear in the prophase of the first cleavage, but in the succeeding telophase a single nucleolus generally appears in each half of each daughter-nucleus. The same is true of the succeeding cleavages, so that each nucleus throughout the cleavage usually has two nucleoli in the telophase or early resting stage.

Beard (1902) described a double structure of the resting stages of the nuclei of the early germ cells of *Raja batis*; these were not traced earlier than a late gastrula stage, but influenced by the findings of Häcker and Rückert, Beard did not hesitate to interpret the double nuclei as consisting of distinct maternal and paternal halves.

Jenkinson (1904) gives some interesting figures of the fertilization and first cleavage stages of *Axolotl*. The germ nuclei meet without fusion, and the chromosomes appear separately in each pronucleus while the nuclear membranes are still intact. In some cases at least, these two chromosome groups remain distinct in the equatorial plate after the dissolution of the nuclear membranes. Scant attention is paid to these features in the text of Jenkinson's paper, which is concerned with other matters, but the author states that he has found two distinct sets of chromosomes in some preparations of the fertilization spindle of *Triton*.

In 1904 Moenckhaus described the independence of the maternal and paternal chromosome groups in the early cleavage spindles of the hybrids produced by fertilizing the eggs of *Fundulus* with *Menidia* sperm. The difference in the size and shape of the chromosomes of the two species makes the identification of the maternal and paternal chromosomes in the case described a matter of certainty.

Pinney (1918) found that two nucleoli are typically present in the nuclei of normal *Ctenolabrus* blastoderms, and cited evi-

dence supporting the view that in these double nucleoli observed in fishes we are dealing with parental homologues.

Concerning *Cryptobranchus allegheniensis*, the writer believes that the observations recorded in the present paper establish beyond question the complete separation of maternal and paternal germ-nuclei to a late blastula stage at least. The separation is particularly marked during the resting stage of the nuclear cycle, precisely where most investigators working with other species have encountered the greatest difficulty.

The observations thus far cited indicate that in certain forms the individuality of the germ-nuclei during early embryonic development is maintained by complete separation of the nuclear material derived from the egg and the spermatozoön respectively. It has already been indicated that this segregation is not by any means a universal phenomenon. Is it possible that the individuality of the germ-nuclei may be maintained, in all essential respects, in those other cases where there is a mingling of chromatin derived from the two germ-nuclei? Let us first examine the facts that require explanation.

In all cases where the germ-nuclei fuse into a single vesicular nucleus before the formation of the first cleavage spindle, mingling of maternal and paternal chromosomes may be expected. Sax (1918) has recently described two cases in flowering plants, *Fritillaria* and *Triticum*. In *Fritillaria* the germ-nuclei usually unite while in the resting condition, although occasionally they are in the spireme stage at the time of fusion. The presence of a single spireme in the zygote could not be demonstrated. In *Triticum* the sperm-nucleus is small and almost homogeneous in structure even while in contact with the egg nucleus. The sperm-nucleus enters the egg-nucleus and there forms a separate compact spireme; at the same time the spireme of the egg-nucleus is formed. In both *Fritillaria* and *Triticum* the maternal and paternal chromosomes are formed independently, but they are not found in separate groups.

Moenckhaus (1904) found that after the first few cleavage divisions in his hybrid teleost eggs the chromosomes of maternal and paternal origin mingled indiscriminately upon the equatorial

plate, and the observations of Morris (1914) are in agreement.

Metz (1916) has found in the diptera a pairing of homologous chromosomes, simulating synapsis, which occurs in all tissues, somatic as well as germinal; this association of maternal with paternal chromosomes was found in late cleavage and during all later stages of embryonic development. From the following statement by Overton (1909) we have a suggestion of a similar occurrence in plants, though attention is directed more particularly to the matter of genetic continuity of individual chromosomes: "In the somatic nuclei (of certain plants) the chromosomes are represented during rest by definite visible bodies, the pro-chromosomes, which are arranged in parallel pairs, with apparent linin intervals. These heterogeneous spirems, the homologous portions of which have early become associated in pairs, probably remain distinct throughout the life-history of the sporophyte."

In spite of the mingling and even paired association of maternal and paternal chromosomes, there are reasons for believing that the two kinds of chromosomes maintain their independence until gametogenesis. This leads us to a consideration of the doctrine of the genetic continuity of individual chromosomes, which goes further than the principle of duality of the embryonic nuclei, but confirms it as a universal law.

The remarkable constancy in the number of chromosomes throughout the cells of a given organism and species has long been known, and affords important evidence for the view that the chromosomes are persistent as individual structures. To be sure, it sometimes happens in mitosis that one or more chromosomes belonging to one daughter-group, accidentally become included with the other group so that one of the daughter-nuclei has fewer, the other more, than the normal somatic number; but such an occurrence is very exceptional, and in subsequent divisions of these cells the number of chromosomes appearing is not the normal, but the increased or diminished number (Boveri, 1890). Whatever the number of chromosomes entering into the formation of a resting nucleus, the same number afterwards issues from it.

In 1883 Van Beneden showed that in *Ascaris* the spermatozoön brings in just as many chromosomes as are contained in the egg. As a result of a careful study of mitosis in epithelial cells of the salamander, Rabl (1885) concluded that the chromosomes do not lose their individuality at the close of division, but persist in the chromatic reticulum of the resting nucleus. Boveri (1887 and 1888) supported Rabl's hypothesis on the ground of his own studies and those of Van Beneden on the early stages of *Ascaris*. Boveri demonstrated in *Ascaris* that in the formation of the spireme the chromosomes reappear in the same position as those which entered into the formation of the reticulum, precisely as Rabl had maintained. As the long chromosomes diverge, their free ends are always turned toward the equator of the spindle, and upon the reconstruction of the daughter-nuclei these ends give rise to corresponding lobes of the nucleus, which persist throughout the resting stage. At the succeeding division the chromosomes reappear exactly in the same position, their ends lying in the nuclear lobes as before. These observations were afterwards confirmed by Herla (1893), and more recently Sutton (1902) has observed practically the same thing in *Brachystola magna*. Boveri (1891) concluded that the chromosomes must be regarded as individuals that have an independent existence in the cell, and expressed his belief that "we may identify every chromatic element arising from a resting nucleus with a definite element that entered into the formation of that nucleus, from which the remarkable conclusion follows that in all cells derived in the regular course of division from the fertilized egg, one half of the chromosomes are of strictly paternal origin, the other half of maternal."

Herla (1893) and Zoja (1895) have shown that if in *Ascaris megalcephala* the egg of variety *bivalens*, having two chromosomes, be fertilized with the spermatozoön of variety *univalens*, having one chromosome, the three chromosomes reappear at each cleavage, at least as far as the twelve-cell stage; and according to Zoja, the paternal chromosome is distinguishable from the two maternal at each step by its smaller size. "We have thus what must be reckoned as more than a possibility, that every cell

of the body of the child may receive from each parent not only half of its chromatin substance, but one half of its chromosomes, as distinct and individual descendants of those of its parents" (Wilson, 1900).

Boveri (1909) found in *Ascaris* that in sister cells preparing for division the configuration of the groups of chromosomes is the same. The similarity of the sister cells is explicable on the view that the chromosomes retain during the resting stage the same shape and size and relative location that they had at the end of the preceding division. In cells of these same embryos that are not sister cells, a great variety of arrangements of the chromosomes is found, and no two arrangements are so nearly alike as are those found in sister cells.

Other evidence for the continuity of individual chromosomes is derived from those cases where the reconstruction of the resting nucleus takes place through the metamorphosis of each chromosome into a hollow vesicle, and the aggregation or fusion of these chromosomal vesicles to form a single nucleus. Such, to be sure, is not the only type of telophase (Wilson, 1900, p. 71), but it takes place in many segmenting ova and in some spermatogonia. According to Conklin (1902), in the late stages of mitosis of the segmenting egg of *Crepidula* the chromosomes enlarge to form vesicles and these unite into a resting nucleus; the nuclear membrane is composed of the outer walls of the vesicles, while the inner walls stretch through the nucleus as chromatic partitions; the chromosomal vesicles from the egg and sperm nuclei respectively remain distinct longer than those from the same germ nucleus. Vesicular chromosomes have been described in fish eggs by Moenckhaus; the individual vesicles fuse with their neighbors and these larger ones with each other until at last the entire nucleus is simply one great vesicle, which is at first lobed, but later is well rounded. Wenrich (1916) says of *Phrynotettix*: "The spermatogonial divisions showed that each chromosome forms a sac or vesicle in the earlier telophases, and that it expands and becomes diffused within these vesicles; that, although the vesicles appeared to coalesce, there is always a remnant of each chromosome visible in the center of the region occupied by the

vesicle, and that in the prophase the chromatin concentrates about this remnant or core and there forms a spirally coiled thread, which develops into a prophase chromosome." Richards (1917) has shown that in the cleavage and gastrula stages of *Fundulus* and *Coregonus* the chromosomes in the telophase of mitosis enlarge to form vesicles which remain distinct though compactly massed together during the resting stage. The new chromosome arises within a vesicle, through the aggregation of its granules; thus there is genetic continuity of individual chromosomes. According to Richards, the polarity of the cell is manifested in the arrangement of the elongated vesicles. When the centrosome divides, the cell acquires a new axis at right angles to the old, in a line connecting the two asters; the chromosomes, when formed, orient themselves with respect to the new axis. Pinney (1918) figures chromosomal vesicles in the telophase of dividing blastomeres of *Ctenolabrus* \times *Fundulus* hybrids. Some observations on chromosomal vesicles in *Cryptobranchus allegheniensis* are included in the present paper. I have examined a few sections of *Coregonus* blastoderms belonging to the collection of the Michigan State Normal College and the material appears to be exceptionally favorable for the study of chromosomal vesicles.

Thus it has been proved that in the fertilized egg one half of the chromosomes are derived from the father and one half from the mother, and that at every division of the egg the chromosomes also divide in such a manner that their progeny are distributed in equal number to all the cells of the egg. Further, there is genetic continuity between each individual chromosome that enters the resting stage and a corresponding chromosome that emerges in preparation for the next division. The conclusion is thus reached that the fertilized egg, and all the cells derived from it, contain a double set of chromosomes, paternal and maternal. This conclusion implies that at every step the respective chromosome groups preserve their independence, however much they are mingled with one another.

This conclusion is further strengthened by observations of individual differences, both morphological and physiological,

between the chromosomes of a given simplex group, and by the behavior of the chromosomes in synapsis. The chromosomes of a single group, maternal or paternal, are not precisely alike, but differ among themselves in size, form and genetic potency (Montgomery, 1901; Sutton, 1902 and 1903; Morgan, 1915). In certain species the size differences are very marked, so that the chromosomes of a single germ-nucleus may be arranged in a graded series; these size differences are constant from one cell-generation to the next, so that individual chromosomes may be identified in successive cell-divisions. In other species where the visible differences are not so marked we have evidence that physiological differences exist, for Bóveri (1907) has shown the strong probability that normal development of the egg is possible only in the presence of at least a single set of qualitatively different chromosomes. Thus the concept of the individuality of the chromosomes has been extended to include not merely the genetic continuity of each particular chromosome, but also its idiosyncrasy or specificity.

In any biparental organism, the duplex chromosome group is composed of two equivalent parental series or simplex groups, in which each individual chromosome is homologous with a very similar chromosome belonging to the other series; in other words, the chromosomes are present in biparental pairs (Montgomery, 1901; Sutton, 1902 and 1903; Wilson, 1912). "In the process known as synapsis, which takes place shortly before the last two cell-divisions concerned in the formation of the germ cells, the chromosomes do in fact unite in pairs, two by two. There is reason to believe that the two members of each pair are respectively of maternal and paternal derivation; and the probability of this view, first stated by Montgomery, has steadily increased." (Wilson, 1913). In one of these two final divisions of the germ cell cycle the double chromosome groups are reduced to single ones in preparation for the subsequent process of fertilization; this reduction is accomplished through the failure of the individual chromosomes to split in the process of mitosis. In this reducing division the two members of a synaptic pair are separated to pass into different daughter cells, but not necessarily

in such a manner that each daughter cell receives exclusively maternal or exclusively paternal chromosomes. The distribution of each pair of homologous chromosomes is entirely independent of that of every other pair, so that in the daughter cells combinations of non-homologous chromosomes occur regardless of their parental origin; each daughter cell usually receives a mixture of maternal and paternal chromosomes in varying proportions. Within the limits of the reduced number, any combination of the chromosomes furnished by the immature germ cell is possible in a daughter cell resulting from the reducing division, save that a single daughter cell cannot receive both members of a synaptic pair. Synapsis is thus explained by the provision which it makes that two homologous chromosomes shall in no case enter the nucleus of a single spermatozoon or mature egg (Sutton, 1902). The principle of random distribution or independent assortment of non-homologous chromosomes has been confirmed by the direct observations of Carothers (1913 and 1917), Wenrich (1915 and 1916) and others. The point that immediately concerns us here is the fact that in the reducing division the individuality of the germ-nuclei is usually lost, for it seldom happens that a single daughter cell, and consequently a single mature gamete, receives exclusively maternal or exclusively paternal chromosomes.

"Synapsis is not a haphazard junction of chromosomes, but an orderly union of elements of maternal and paternal origin, similar in size, in details of form, and probably also in function" (Kellicott, 1913, p. 294). The orderly processes of heredity as they exist today in biparental organisms would be impossible were the integrity of the maternal and paternal chromosome groups not maintained up to the time of synapsis. In *Cryptobranchus allegheniensis* and in some other forms this integrity is manifested at every step by a complete segregation of maternal and paternal chromosome *groups*; in other forms the independence of the chromosome groups is maintained in spite of the mingling of maternal and paternal chromosomes. The persistent individuality of the chromosomes is the important thing, and so long as this is maintained the germ-nuclei exist as actual

entities whether visibly separated or not. The distinction between the two classes of germ-nuclei—those that fuse at the time of fertilization and those that do not—is more apparent than real. The expression “individuality of the germ-nuclei” seems justified in either case, for individuality implies separableness as well as separateness (Conklin, 1916); but some may prefer to use the term “autonomy,” which places the emphasis upon independence. Since attention has become focused upon the chromosomes, the expression “autonomy of the maternal and paternal chromosome groups” has often been used to indicate the duality of nuclear structure.

Where it exists, the separation of maternal and paternal chromatin-complexes into distinct groups within a single nucleus affords a striking exemplification of the deeper and more universal truth that each germ-nucleus is represented in its entirety in every cell, somatic as well as germinal, of a developing organism. In *Cryptobranchus allegheniensis* this dual structure of the nucleus is clearly visible in every cell of the early segmentation stages at least, so that here we have material for an ocular demonstration of a principle long ago foreseen by Huxley (1878), who wrote: “It is conceivable, and indeed probable, that every part of the adult contains molecules derived from both the male and the female parent; and that, regarded as a mass of molecules, the entire organism may be compared to a web of which the warp is derived from the female and the woof from the male.”

SUMMARY.

In the fertilization of the egg of *Cryptobranchus allegheniensis* the germ-nuclei do not fuse, and in the first cleavage mitosis each gives rise to a separate group of chromosomes whose descendants pass separately to the daughter-nuclei.

During the ensuing resting stage each germ-nucleus is represented by a structurally distinct vesicle. The separateness of the germ-nuclei is thus maintained throughout the entire nuclear cycle.

Throughout early cleavage the nuclear divisions are of the same duplex type, and the resting nuclei are always distinctly

double. The genetic continuity of each half of the double nucleus has been clearly traced to an advanced cleavage stage.

During late cleavage and in the early gastrula the nuclei are still typically double, but certain irregularities which tend to disguise the double structure occur with increasing frequency and the segregation of maternal and paternal chromatin cannot always be demonstrated.

The hypothesis of individuality of the germ-nuclei as applied to those species in which there is a mingling of maternal and paternal chromosomes is discussed, and supported by considerations regarding the persistent individuality of the chromosomes.

ZOOLOGICAL LABORATORY,
MICHIGAN STATE NORMAL COLLEGE,
YPSILANTI.

REFERENCES.

Beard, J.

'02 The germ cells. I. *Raja batis*. Zoöl. Jahrb., 16.

Van Beneden, E.

'83 Recherches sur la maturation de l'oeuf, la fécondation et la division cellulaire. Archiv Biol., IV.

Boveri, Th.

'87 Ueber den Anteil des Spermatozoön an der Teilung des Eies. Sitz.-Ber. d. Ges. f. Morph. und Phys. in München, Bd. III., Heft 3.

'88 Zellenstudien, II. Die Befruchtung und Teilung des Eies von *Ascaris megalcephala*. Jena. Zeit., 22.

'90 Zellenstudien, III. Ueber das Verhalten der chromatischen Kernsubstanz bei der Bildung der Richtungkörper und bei der Befruchtung. Jenaische Zeitschrift, 24.

'91 "Befruchtung" in Merkel und Bonnet's Ergebnisse, I., p. 410.

'07 Zellenstudien, VI. Die Entwicklung dispermer Seeigel-Eier. Ein Beitrag zur Befruchtungslehre und zur Theorie des Kernes. Jena. Zeit., 43.

'09 Die Blastomerenkerne von *Ascaris megalcephala* und die Theorie der Chromosomen-Individualität. Archiv Zellf., III.

Carothers, E. Eleanor.

'13 The Mendelian ratio in relation to certain Orthopteran chromosomes. Jour. Morph., Vol. 24.

'17 The segregation and recombination of homologous chromosomes as found in two genera of Acrididae (Orthoptera). Jour. Morph., Vol. 28, No. 2.

Conklin, E. G.

'01 The individuality of the germ-nuclei during the cleavage of *Crepidula*. BIOL. BULL., II.

'02 Karyokinesis and cytokinesis in the maturation, fertilization and cleavage of *Crepidula* and other gasteropods. Journal of the Academy of Natural Sciences, Philadelphia, Vol. 12.

'16 The basis of individuality in organisms, from the standpoint of cytology and embryology. Science, April 14.

Guyer, Michael F.

- '11 Recent progress in some lines of cytology. Trans. Amer. Micros. Soc., April.

Häcker, V.

- '92 Die Eibildung bei *Cyclops* und *Canthocamotus*. Zoöl. Jahrb., Bd. V.
'95 Ueber die Selbständigkeit der väterlichen und mütterlichen Kernbestandtheile während der Embryonalentwicklung von *Cyclops*. Archiv für mikr. Anat., Bd. 46.

Herla, V.

- '93 Étude des variations de la mitose chez l'*Ascaride Megalocephale*. Archiv Biol., XIII.

Hertwig, Oscar

- '75-'78 Beiträge zur Kenntniss der Bildung, Befruchtung und Theilung des Thierischen Eies. Morph. Jahrb., 1-4.

Huxley, T. H.

- '78 Article "Evolution" in Encyclopædia Britannica, VIII., pp. 744-751.

Jenkinson, J. W.

- '04 Observations on the maturation and fertilization of the egg of the Axolotl. Q. J. M. S., Vol. 48.

Kellicott, Wm. E.

- '13 General embryology. Henry Holt and Co.

Lillie, Frank R.

- '16 The history of the fertilization problem. Science, January 14.

Mark, E. L.

- '81 Maturation, fecundation and segmentation of *Limax campestris*. Bull. Mus. Comp. Zoöl., Harvard College, VI.

Metz, Charles W.

- '16 Chromosome studies in the diptera. II. The paired association of the chromosomes in the diptera, and its significance. Jour. Exp. Zoöl., Vol. 21, No. 2.

Moenckhaus, W. J.

- '04 The development of the hybrids between *Fundulus heteroclitus* and *Menidia notatus*, with especial reference to the behavior of maternal and paternal chromosomes. Jour. Anat., Vol. 3, pp. 29-65.

Montgomery, Thos. H.

- '01 A study of the chromosomes of the germ cells of the metazoa. Trans. Amer. Phil. Soc., XX.
'04 The main facts in regard to the cellular basis of heredity. Proc. Amer. Phil. Soc., Vol. XLIII, No. 175.

Morgan, Sturtevant, Muller, Bridges

- '15 The mechanism of Mendelian heredity. Henry Holt and Co.

Morris, Margaret

- '14 The behavior of the chromatin in hybrids between *Fundulus* and *Ctenolabrus*. Jour. Exp. Zoöl., Vol. 16.

Parmenter, Charles Leroy.

- '19 Chromosome numbers and pairs in the somatic mitoses of *Amblystoma tigrinum*. Jour. Morphol., Vol. 33, No. 1, Dec. 20.

Overton, James Bertram

- '09 On the organization of the nuclei in the pollen mother-cells of certain plants, with especial reference to the permanence of the chromosomes. *Annals of Botany*, Vol. 23, No. 89, January.

Pinney, Edith

- '18 A study of the relation of the behavior of the chromatin to development and heredity in teleost hybrids. *Jour. Morph.*, Vol. 31, No. 2, September.

Rabl, C.

- '85 Ueber Zelltheilung. *Morph. Jahrb.*, X.

Richards, A.

- '17 The history of the chromosomal vesicles in *Fundulus* and the theory of the genetic continuity of the chromosomes. *BIOL. BULL.*, Vol. XXXII., No. 4, April.

Rückert, J.

- '95 Ueber das Selbständigbleiben der väterlichen und mütterlichen Kernsubstanz während der ersten Entwicklung des befruchteten *Cyclops*-Eies. *Archiv für mikr. Anat.*, 45.

Sax, Karl

- '18 The behavior of the chromosomes in fertilization. *Genetics*, Vol. 3, No. 4, July.

Smith, Bertram G.

- '12 The embryology of *Cryptobranchus allegheniensis*. Part I: Introduction; the history of the egg before cleavage. *Jour. Morph.*, Vol. 23, No. 1, March. Part II: General embryonic and larval development, with special reference to external features. *Jour. Morph.*, Vol. 23, No. 3, September.

Sutton, W. S.

- '02 On the morphology of the chromosome groups in *Brachystola magna*. *BIOL. BULL.*, Vol. IV.
- '03 The chromosomes in heredity. *BIOL. BULL.*, Vol. IV.

Wenrich, D. H.

- '15 Synapsis and the individuality of the chromosomes. *Science*, N. S., Vol. 41, No. 1055.
- '16 The spermatogenesis of *Phrynotettix magnus*, with special reference to synapsis and the individuality of the chromosomes. *Bull. Mus. Comp. Zoöl.*, Harvard College, Vol. 60, No. 3.

Wilson, E. B.

- '00 The cell in development and inheritance. The Macmillan Co.
- '12 Studies on chromosomes, VIII. Observations on the maturation phenomena in certain hemiptera and other forms, with consideration of synapsis and reduction. *Jour. Exp. Zoöl.*, Vol. 13.
- '13 Heredity and microscopical research. *Science*, May 30.

Zoja, Raffaello

- '95 Sulla indipendenza della cromatina paterna e materna nel nucleo delle cellule embrionali. *Anat. Anz.*, Bd. XI., No. 10.

EXPLANATION OF PLATE I.


Cryptobranchus allegheniensis.

FIG. 1. Portion of a meridional section of a fertilized egg, showing the meeting of the sperm-nucleus and the egg-nucleus. Each germ-nucleus extends through four sections, of which an inner one is shown in the figure. The section passes midway between the two asters, which lie on opposite sides of the nuclei in a line at right angles to the plane of the section. The egg was fixed with bichromate-acetic formalin and stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 2. Meridional section of an egg killed twenty-two and one half hours after fertilization, showing the resting germ-nuclei. Each germ-nucleus extends through five sections, of which the middle one is here shown. The section passes midway between the two asters, which lie close to the apposed surfaces of the two nuclei in a line oblique to the plane of the section. Fixed with bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.

FIG. 3. Horizontal section through the resting germ-nuclei of an egg killed twenty-one and one half hours after fertilization. Nearly half of each nucleus is left in an adjacent section. Two asters are shown in the figure. Fixed with bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 4. Horizontal section through the resting germ-nuclei of an egg killed twenty-two and three fourths hours after fertilization. Each germ-nucleus extends through three sections, of which the middle one is here shown. Two asters are shown in the figure. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.




1



2



3



4

EXPLANATION OF PLATE II


Cryptobranchus allegheniensis.

FIG. 5. Horizontal section of a fertilized egg showing the germ-nuclei preparing for the first cleavage mitosis. The germ-nucleus shown at the right is in a slightly more advanced phase than the other. A portion of each nucleus is left in an adjacent section. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 6. Horizontal section of an egg killed twenty-two and one half hours after fertilization, showing the germ-nuclei preparing for the first cleavage mitosis. The germ-nucleus shown at the right is in a more advanced phase than the other. About half of each nucleus is left in an adjacent section. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.

FIG. 7. Horizontal section through the first cleavage spindle (late prophase). The chromosomes are segregated into two groups, of maternal and paternal origin respectively. A very small portion of each group of chromosomes is left in each of the adjacent sections. Bichromate-acetic-formalin fixation; stained with iron-hæmatoxylin. $\times 400$.


FIG. 8. Horizontal section of an egg killed twenty-six hours after fertilization, showing a late stage (early telophase) of the first nuclear division. In the distribution of chromatin to the daughter-nuclei, the segregation of maternal and paternal chromatin is maintained. A very small portion of the chromatin of each daughter-nucleus is left in an adjacent section. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric acid. $\times 400$.



5



6



7



8

EXPLANATION OF PLATE III.

Cryptobranchus allegheniensis.


FIGS. 9 AND 10. Two successive vertical sections through a single daughter-nucleus of the first cleavage mitosis. The sections are nearly meridional, and pass very close to the other daughter-nucleus, hence at right angles to the plane of the first cleavage furrow which is not yet formed. The nucleus shown is almost entirely confined to the two sections. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric acid. $\times 400$.

FIG. 11. Vertical section through a single daughter-nucleus of the first cleavage mitosis. The section is nearly meridional, and passes very close to the other daughter-nucleus, hence at right angles to the plane of the first cleavage furrow which is not yet formed. A considerable part of each nuclear vesicle is left in an adjacent section, and comparison with these other portions shows that the plane of contact of the two nuclear vesicles is oblique to the plane of the section. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 12. Vertical section showing a single daughter-nucleus of the first cleavage mitosis, taken at right angles to the plane of the first cleavage furrow which is not yet formed. The entire double nucleus is confined to this section. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.

FIG. 13. Vertical section, parallel to the newly-formed first cleavage furrow, through a single daughter-nucleus of the first cleavage division. Part of the nucleus is left in an adjacent section, where it is likewise distinctly double. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 14. Vertical section, parallel to the first cleavage furrow, through the other daughter-nucleus of the egg used for Fig. 13. Part of the nucleus is left in an adjacent section, where the two nuclear vesicles are more closely united. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.



9




10




11



12



13



14

EXPLANATION OF PLATE IV.

Cryptobranchus allegheniensis.


FIGS. 15 AND 16. Two successive horizontal sections showing a single nucleus rotating into position for the second cleavage mitosis; almost the entire nucleus is confined to these two sections. The newly-formed first cleavage furrow extends in a direction parallel to a line connecting the two asters. Fixed in bichromate-acetic-formalin; stained with iron-hæmatoxylin. $\times 400$.

FIG. 17. Horizontal section through a single nucleus in position for the second cleavage mitosis; the greater part of the nucleus lies in this section. The newly-formed first cleavage furrow extends in a direction parallel to a line connecting the two asters. Fixed in Zenker's fluid; stained with iron-hæmatoxylin. $\times 400$.


FIG. 18. Horizontal section through the other nucleus of the egg containing the nucleus shown in Fig. 17. Orientation as in the preceding figure. Stained with iron-hæmatoxylin. $\times 400$.

FIG. 19. Horizontal section showing a single nucleus in position for the second cleavage mitosis; the entire nucleus lies in this section. The newly-formed first cleavage furrow extends in a direction parallel to a line connecting the two asters. Bichromate-acetic-formalin fixation; stained with borax-carmine Lyons-blue picric-acid. $\times 400$.


FIG. 20. Horizontal section through the other nucleus of the egg used for Fig. 19. A small part of one nuclear vesicle, and the greater part of the other, lies in an adjacent section. Orientation as in the preceding figure. Stained with borax-carmine Lyons-blue picric-acid. $\times 400$.




15




16




17



18



19



20


EXPLANATION OF PLATE V.

Cryptobranchus allegheniensis.


FIGS. 21 AND 22. Two successive meridional sections through a single nucleus preparing for the second cleavage mitosis. The sections extend at right angles to the first cleavage furrow, and close to the other cleavage-nucleus. A considerable part of the nucleus here shown is left in an adjacent section where it is not so clearly separated into two nuclear vesicles. Fixed in bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.

FIG. 23. Horizontal section through a single nucleus preparing for the second cleavage mitosis; the nucleus is almost entirely confined to this section. The first cleavage furrow extends in a direction nearly parallel to a line connecting the two asters. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIGS. 24, 25 AND 26. Three successive horizontal sections through a single nucleus entering upon the second cleavage mitosis; all but a very small portion of the nucleus is included in these sections. The first cleavage furrow extends in a direction nearly parallel to a line connecting the two asters. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.




21




22




23



24



25



26

EXPLANATION OF PLATE VI.

Cryptobranchus allgeheniensis.

FIG. 27. Horizontal section through a nucleus in a late prophase of the second cleavage mitosis. A small portion of each group of chromosomes is left in each of the adjacent sections. The axis of the spindle lies in a direction nearly parallel to the plane of first cleavage. Fixed with bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 28. Horizontal section through one of the daughter-nuclei in a late stage (early telophase) of a second cleavage mitosis. Orientation as in the preceding figure; the second cleavage furrow is not yet formed. Fixed in Zenker's fluid; stained with iron-hæmatoxylin. $\times 400$.

FIG. 29. Horizontal section through one of the daughter-nuclei of a second cleavage mitosis; early resting stage. Part of the nucleus is left in an adjacent section where it is likewise distinctly double. The plane of apposition of the two nuclear vesicles is parallel to the first cleavage furrow, and at right angles to the second cleavage furrow which is beginning to form. Bichromate-acetic-formalin fixation; stained with iron-hæmatoxylin. $\times 400$.


FIG. 30. Horizontal section through one of the daughter-nuclei of a second cleavage mitosis; resting stage. A portion of the nucleus is left in each of the adjacent sections where it is likewise distinctly double. The plane of apposition of the two nuclear vesicles is parallel to the plane of first cleavage, and at right angles to the beginning second cleavage furrow. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.

FIG. 31. Vertical section through one of the daughter-nuclei of a second cleavage mitosis; resting stage. The section is taken parallel to the plane of the second cleavage furrow which has just begun to form. Part of the nucleus is left in an adjacent section, where it is likewise distinctly double. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 32. Horizontal section through a nucleus rotating into position for the third cleavage mitosis; almost the entire nucleus is confined to this section. The second cleavage furrows are present and extend nearly to the equator of the egg. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.




27




28




29



30



31



32

EXPLANATION OF PLATE VII.

Cryptobranchus allegheniensis.

FIG. 33. Horizontal section through a resting nucleus in position for the third cleavage mitosis. The plane of apposition of the two nuclear vesicles is nearly parallel to the *second* cleavage furrows. A considerable part of the nucleus is left in an adjacent section, where it is likewise distinctly double. Fixed with bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 34. Horizontal section through a nucleus in position for the third cleavage mitosis. A line connecting the two asters is nearly parallel to the second cleavage furrow. Part of the nucleus is left in an adjacent section, where it is likewise distinctly double. Fixed with bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.

FIG. 35. Vertical section through a nucleus entering upon the third cleavage mitosis. Part of the nucleus is left in each of the adjacent sections, where it is likewise distinctly double. Fixed with bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 36. Horizontal section through a nucleus in the late prophase of the third cleavage mitosis. A line connecting the two asters extends nearly at right angles to the first cleavage furrow. Fixed with bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.

FIG. 37. Horizontal section through a daughter-nucleus of a third cleavage mitosis. The two nuclear vesicles remain almost in the position in which they were formed, but two asters are present and these have taken up a position preparatory to the fourth cleavage mitosis. A line connecting the two asters extends in a direction almost parallel to the adjacent newly-formed third cleavage furrow which is vertical and intersects a second furrow almost at right angles. Fixed in bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 38. Horizontal section through a nucleus in position for the fourth cleavage mitosis. A line connecting the two asters extends parallel to the adjacent third cleavage furrow and almost parallel to the first furrow. Fixed in bichromate-acetic-formalin; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.




33




34




35



36



37



38

EXPLANATION OF PLATE VIII.

Cryptobranchus allegheniensis.

FIGS. 39 AND 40. Two successive vertical sections through a nucleus preparing for the fourth cleavage mitosis; almost the entire nucleus lies in these two sections. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.


FIG. 41. Another nucleus found in the section used for Fig. 39; early prophase of the fourth cleavage mitosis. $\times 400$.

FIG. 42. Vertical section through a nucleus in the metaphase of the fourth cleavage mitosis. The fourth cleavage furrow has not yet been formed. Bichromate-acetic-formalin fixation; stained with iron-hæmatoxylin. $\times 400$.


FIGS. 43 AND 44. Two successive horizontal sections through a newly-formed daughter-nucleus of a fourth cleavage mitosis. The fourth cleavage furrow is newly formed. Bichromate-acetic-formalin fixation; stained with borax-carminc Lyons-blue picric-acid. $\times 400$.




39




40



41



42



43




44

EXPLANATION OF PLATE IX.


Cryptobranchus allegheniensis.

FIG. 45. Diagrams illustrating the orientation of the germ-nuclei during the first four cleavage divisions. All the diagrams represent horizontal sections, and the germ-nuclei are assumed to lie always in the same horizontal plane. The size of the nucleus is exaggerated in proportion to the size of the egg. Dotted lines represent beginning cleavage furrows; solid straight lines represent cleavage furrows which have cut to the level of the nuclei. The number of chromosomes formed from each germ-nucleus is assumed to be six, and this is probably the correct number.


- A. Germ-nuclei and asters in position for the first cleavage mitosis.
- B. First cleavage mitosis.
- C. Newly-formed daughter-nuclei of the first cleavage mitosis.
- D. Nuclear vesicles and asters in position for the second cleavage mitoses.
- E. Second cleavage mitoses.
- F. Newly-formed daughter-nuclei of the second cleavage mitoses.
- G. Nuclear vesicles and asters in position for the third cleavage mitoses.
- H. Third cleavage mitoses.
- I. Newly-formed daughter-nuclei of the third cleavage mitoses.
- J. Nuclear vesicles and asters in position for the fourth cleavage mitoses.
- K. Fourth cleavage mitoses.
- L. Newly-formed daughter-nuclei of the fourth cleavage mitoses.




A




B




C




D




E




F




G




H




I



J



K



L

BIOLOGICAL BULLETIN

THE NATURE OF THE FERTILIZATION MEMBRANE OF *ASTERIAS* AND *ARBACIA* EGGS.¹

WALTER E. GARREY.

It is generally believed that the fertilization membrane of an echinoderm egg is a distinct structure formed by the egg, and separated from it by the liquid filled perivitelline space. Views quite at variance with this current one have been advanced in recent communications; these we believe are not justified by the experimental evidence which is to be considered below.

I. Elder,² Kite³ and McClendon⁴ have separately maintained that the egg jelly is essential to the formation of the fertilization membrane and in one way or another enters into its composition. Harvey⁵ has made this question one of special study and presents convincing experimental evidence to show that the egg jelly is not essential. He was able by repeated shaking and continued washing with sea water, to remove this jelly so completely that not a trace could be detected when the eggs were examined in India ink suspensions. Fertilizations of such eggs caused the formation of fertilization membranes, which however may be more tenuous than normally, probably due to the fact that the jelly facilitates the membrane formation by acting as a mechanical block to the diffusion of the membrane forming colloid ("membranogen") which was thus retained in more concentrated solution and consequently formed a denser membrane.

J. Loeb⁶ found that hydrochloric acid would dissolve the jelly

¹ From the Physiological Laboratory of Tulane University, New Orleans, and the Marine Biological Laboratory, Woods Hole.

² Elder, *Arch. f. Entwickl.*, 1913, XXXV., 195.

³ Kite, G. L., *Science*, 1912, N.S., XXXVI., 562.

⁴ McClendon, J. F., *Internat. Zeitschr. f. physik. chem. Biol.*, 1914, I., 163.

⁵ Harvey, E. N., *BIOL. BULL.*, 1914, XXVII., 237.

⁶ Loeb, J., *Science*, 1914, N.S., XL., 318.

from the eggs of *Strongylocentrotus purpuratus* and that after subsequent treatment with sodium hydroxide and calcium chloride, starfish sperm caused the formation of fertilization membranes and induced development. F. R. Lillie¹ found that *Arbacia* "eggs are fertilizable" after complete removal of the jelly by Loeb's method. Loeb and Kupelwieser² caused the formation of fertilization membranes by treating eggs with butyric acid; these were freed from every vestige of membrane and jelly was removed by shaking, yet upon insemination they were able to form new fertilization membranes. Moore³ noted that egg fragments, produced by shaking, formed membranes upon entrance of the spermatozoön. These observations indicate that the egg jelly is not essential for the formation of the fertilization membrane.

II. The facts considered in the preceding section make us doubt the validity of the hypothesis advanced by McClendon that the fertilization membrane is formed by the process of precipitation due to the contact of two colloids carrying opposite electric charges; viz., the egg jelly which McClendon states is electro-negative and another colloid (membranogen) derived from the egg which he found to be electropositive. This view is rendered quite untenable when we consider the fact that fertilization membranes are formed about echinoderm eggs by Loeb's well-known methods of inducing artificial parthenogenesis by treatment with weak fatty acids. These acids act only after penetrating the egg-jelly in the course of which event they impart to that colloid the positive electric charge of the dominant hydrogen ion; they therefore have the same charge as the colloids of the egg and the requisite conditions for precipitation of colloids do not exist.

It has furthermore been shown that in the chemical fertilization of *Asterias* eggs either acids or alkalis may be used to induce membrane formation. The electric charge of the jelly is positive when acids are used and negative with alkalis; obviously the charge is not of opposite sign to that of the egg substance (membranogen) in both cases.

¹ Lillie, F. R., *Biol. Bull.*, 1915, XXVIII., 24.

² Kupelwieser, H., *Arch. f. Entw.*, 1909, XXVII., 434.

³ Moore, A. R., *Univ. California Pub.*, 1912, IV., 89.

Upon treatment with acids the egg jelly swells perceptibly and also removes the acid from solution either by adsorbing it or by chemically uniting with it. In either event the jelly becomes electropositive. The fixation of the acid is shown in the following experiments.

Ripe eggs from *Asterias* ovaries are shed in sea water and allowed to settle to the bottom of a graduated cylinder. The supernatant liquid is decanted and enough fresh sea water added to the mass of eggs to equal three times their volume. The cylinder is actively shaken and the eggs again are allowed to settle. The clear supernatant fluid containing a considerable amount of the jelly is now decanted off and an equal volume of $N/500$ butyric acid added. Fresh *Asterias* egg do not form fertilization membranes in this acid-jelly mixture but do so in a control solution of $N/1,000$ butyric acid in sea water. On the other hand fertilization membranes are formed at once in the acid-jelly mixture after the further addition of an equal volume of $N/500$ butyric acid sea water, thus indicating clearly that the initial effect of the jelly is to remove the acid from solution and that membrane formation is induced only when acid is present in excess of the amount necessary to saturate the jelly. Neglect in the observance of this precaution doubtless has been the cause of many failures to obtain good artificial fertilization by this method in case large masses of eggs have been added to small amounts of acid sea water. Loeb¹ cautioned against such a loose procedure. Under such conditions the relatively large amount of adherent jelly fixes the acid and prevents it from acting on the eggs.

III. The formation of fertilization membranes after the complete removal of the jelly indicates the origin of this structure from the egg. The following experiment furnishes a simple but equally conclusive mechanical demonstration of this fact. Fertilization of the eggs of *Asterias* either by sperm or by artificial means such as acid or heat ($30-33^{\circ}$ C.) is possible not only after maturation but also earlier, when the nuclear membrane about the germinal vesicle just begins to fade (De Lage²); in fact I have

¹ Loeb, J., "Artificial Parthenogenesis" (Chicago), 1913, 69.

² DeLage, Y., *Arch. de zool. expér. et gen.*, ser. 3, IX., 285.

often observed that fertilization is possible even before the nuclear membrane begins to fade. After both polar bodies were formed I found that fertilization by any means caused the formation of the membrane between them and the egg surface, from which they were lifted and pushed away as the membrane moved outward. They are seen to lie in the saucer-like depression in the outer surface of the membrane, which results from the resistance offered by the egg-jelly to the pressure developed in the perivitelline space. After staining with dilute methylene blue, examination removes all doubt that the polar bodies are entirely outside the membrane.

By so timing the fertilization that only the first polar body has been formed, it will be pushed away from the egg outside the fertilization membrane while the second polar body, being formed much later, remains at the surface of the egg, and therefore inside the perivitelline space. Similarly by fertilizing before maturation the fertilization membrane will have been formed and will have left the surface of the egg before the polar bodies are extruded. These remain at the surface of the egg and therefore within the perivitelline space. By manipulation they may be shaken free from the surface of the egg and moved about freely.

IV. Kite¹ has stated that the "so-called fertilization membrane of the egg of *Arbacia* consists of three parts, viz., the inner layer of the egg jelly which has undergone a change in refraction index, the swollen vitelline membrane, and the thin highly refractive surface layer of the cytoplasm. This hyaline layer is still very adherent to the vitelline membrane." Kite thus conceives the vitelline membrane to remain attached to the egg, to become swollen and edematous and to fill completely the entire space between the egg and the jelly which in turn is altered in refractive power to appear like a separate membrane. That the egg jelly is non-essential has been shown. The other features of Kite's conception can likewise be demonstrated to be erroneous both for *Arbacia* and *Asterias*. In these forms what appears to be the fertilization membrane is far more than the outer refractive part of an invisible thick layer of "gel" formed by a swelling of the vitelline membrane. It is a true, thin membrane formed

¹ Kite, G. L., *Science*, 1912, N.S., XXXVI., 562.

by the egg and separated from its surface by a liquid filled space as is indicated by the following considerations:

1. This new structure has entirely different permeability from that of the surfaces of either the fertilized or unfertilized eggs which are very slightly permeable to neutral salts, so that the eggs are plasmolyzed and crenated by hypertonic saline solutions. Salts however must penetrate the fertilization membrane freely to effect this change in the egg and that they do so is further shown by the fact that the fertilization membrane retains its round contour and distended condition when placed in the hypertonic or hypo-tonic salt solutions. The differences in permeability thus far demonstrated speak against the origin of the membrane from a pre-existing structure on the surface of the egg.

2. Kite's description is proven incorrect by the demonstration of the fact that the fertilization membrane is separated from the surface of the egg by a space filled with liquid and not occupied by a swollen gelatinous vitelline membrane continuous with the surface of the egg on the inside and with the egg jelly on the outside. Dr. Robt. Chambers by removing this liquid from this space for me by the use of his micro-pipette has produced a collapse of the membrane. Allusion has already been made to the fact that polar bodies extruded after the formation of the fertilization membrane may be separated from the egg surface by manipulation and are then free to assume various positions in the space; this could be possible only in liquid but certainly not if the space was occupied by any sort of a "gel."

Professor J. Loeb has shown that the fertilization membrane of echinoderms is impermeable to such colloids as egg white, shark's serum, and even tannic acid. These materials, if added to sea water, cause the thin membrane to crumple onto the surface of the enclosed egg by the osmotic extraction of water from the intervening space; but the membrane regains its original contour when replaced in sea water. This latter fact suggested to Loeb the probability that the space contains some colloid in solution secreted by the egg at the time the fertilization membrane is formed. To this colloid the membrane is impermeable; it therefore exerts its osmotic pressure in excess of that due to the salts of the sea water, serves to raise the membrane from the surface of the egg and to keep it distended.

Experiments by the writer have shown that when fertilized eggs of *Asterias* or *Arbacia* are placed in a two per cent. solution of Witte's peptone in sea water the fertilization membrane crumples instantly as in Loeb's experiments, but that when allowed to remain in the solution the membranes are again distended, indicating that they are slowly permeated by the albumoses in this preparation. Such results can only be obtained where a semi-permeable membrane, one permeable to salts but impermeable to colloids, encloses a space containing a colloidal solution.

4. Still other facts show that the space between the fertilization membrane and the egg surface is filled with a fluid and not with a gel. If *Asterias* eggs are heated to 33° C. they form fertilization membranes in from three to five minutes. By further warming for fifteen minutes they show slow amœboid movements after they have been returned to sea water at 20° C. They may migrate about in the space enclosed by the thin fertilization membrane and up close to that structure at any point. They may throw out long streamer like pseudopodia which meet with no check to their progress anywhere in the space till they reach the thin confining fertilization membrane. This over-heating process may lead to droplet formation and ultimately to partial or complete disintegration of the egg; the débris thus formed is scattered throughout the space and is never held away from the thin fertilization membrane by any gelatinous structure. Similarly if the fertilized eggs of *Asterias* are subjected to the action of diluted sea water they may be caused to swell until each presses tightly against the confining fertilization membrane and completely fills the space. Cytolysis will result in sea water which has been sufficiently diluted and in this case, as in that of cytolysis caused by prolonged warming, the granular material completely fills the perivitelline space which may be more than twice the original diameter of the egg. When extra-ovates rupture through the fertilization membrane after treatment with hypotonic solutions, the egg is drawn over tightly against the inner surface of the membrane at the point of rupture, the thinness of the membrane being evident at the constricted neck between the extra-ovate and the main egg mass.

Again, in the formation of the blastula the cells range up close to the inner surface of the fertilization membrane and completely obliterate the interior space giving a further indication that neither a thickened vitelline membrane nor any "gel" is there.

These demonstrations may be made immediately after fertilization so that there is no necessity to assume that the vitelline membrane or a gel in the perivitelline space has had time to undergo liquefaction.

The above facts and others which might be referred to all indicate the fluid character of the contents of the space between the fertilization membrane and the egg surface and serve to substantiate the conception of the fertilization membrane stated in the opening paragraph of this communication. In the case of both *Arbacia* and *Asterias* the fertilization membrane is a structure which arises at the surface of the egg and is subsequently lifted from it and distended by fluid. Depending upon the time of fertilization of *Asterias* eggs relative to the formation of the polar bodies these structures may be outside the fertilization membrane, inside the perivitelline space or the first outside and the second within the fertilization membrane.

Heilbrunn¹ has reviewed the subject of membrane production by *Arbacia* eggs under the influence of various chemicals. He concludes with Herbst and others that in this form the membrane exists preformed on the surface of the unfertilized egg and that fertilizing agents merely cause its elevation. He gives evidence in support of the view that this elevation is the result of lowering of surface tension and swelling of proteins beneath the membrane. In the light of our findings it is clear that the fluid character of the contents of the intra-membranal space and the progressive increase in its bulk as the membrane is lifted away from the egg indicate that the colloids are in solution and that the process involves the osmotic attraction of water.

It is not yet demonstrated that a membrane exists preformed on the surface of unfertilized *Asterias* egg and the formation of a membrane on a fragment of *Arbacia* egg as described by Moore (*loc. cit.*) are facts which speak, in these instances, for the formation of the fertilization membrane *de novo*.

¹ Heilbrunn, Lewis V., BIOL. BULL., 1913, XXIV, p. 343.

THE EFFECT OF ULTRA-VIOLET LIGHT RAYS UPON THE DEVELOPMENT OF THE FROG'S EGG.

W. M. BALDWIN,

DEPARTMENT OF ANATOMY, UNION UNIVERSITY (ALBANY MEDICAL COLLEGE).

II THE ARTIFICIAL PRODUCTION OF FOLDED (U-SHAPED) EMBRYOS.

This paper is the second of a series dealing with an investigation of the nature and mechanism of causation of certain abnormal, developmental results obtained by raying certain restricted areas of the fertilized ovum of the frog by means of ultra-violet light rays. Since the first paper dealt with the production of a single type-defect, spina bifida, the possibility of production of another constant and fixed experimental result, a folded or U-shaped embryo, justifies the separate consideration of that result in this contribution. Once again the value of the physical method of attack upon the complex problem of developmental reaction is demonstrated. In this problem we are concerned with a readily controllable and constant, causative agent on the one hand and a uniform embryological defect on the other. This one technical condition permits of a more thorough analytical inquiry into the mechanism of production of this developmental defect. In addition to this consideration, however, as is to be expected the experiment gives further insight into the genetic constitution of the fertilized, undivided ovum, and, more specifically, helps in the identification and location of the organ forming substances and anlagen. The apparatus used for the experiment was that detailed in the preceding paper. Some of the work was done, however, with an apparatus for the use of which I am indebted to Mr. W. S. Andrews, of the General Electric Company at Schenectady. With both forms of apparatus, the surface area concentration of the rays was increased through the use of convex quartz lenses.

The eggs used were those of the common species of frogs found

around Ithaca and also Schenectady, N. Y. These were collected early in the morning, as soon as possible after laying. They were brought into the laboratory, divested of their jelly, and rayed while still in the undivided or two-celled stage. A thin tin-foil diaphragm perforated by a hole three tenths of a millimeter in diameter shielded most of the surface area of the eggs from the rays but permitted a relatively small portion of that area to be influenced. After raying, the eggs were transferred to a jar containing 1,000 c.c. of tap water in which they were permitted to develop, the water being changed frequently.

Physicists have definitely established the fact that ultra-violet light rays possess a very slight penetration capacity. We are justified, hence, in assuming that the chemical alterations produced by the rays acting under these conditions were restricted to a superficial layer of protoplasm of the egg in the restricted area mentioned. The interpretations presented in this paper are made, accordingly, upon that basis.

The surface area of the egg subjected to examination in this experiment is best demonstrated by a reference to Fig. 1. A




FIG. 1.

two-cell ovum is shown with the animal pole uppermost. The broken line marks out the area investigated. As is to be seen this extends from the region of the equator up towards the animal pole but is exclusive both of the pole itself and of a narrow median strip which connects the latter with the equator. As can be seen this region is exclusive of the gray crescent. Different portions of this area were rayed successively with the three tenths millimeter beam of light. As a result, when development

was permitted to progress without further interference, the long axis of the trunk of the embryos was invariably bent towards one or the other side. The level of this bending or flexure of the trunk depended directly upon the location of the area influenced by the light.

Since the jelly which envelops the eggs had been found in the spina bifida experiments to be impervious to ultra-violet light rays, care was taken to remove it prior to the experiments. As a result a marked increase in the percentage of positive results followed the raying. Where, as was the case more particularly with green frog's eggs, the jelly could be removed almost completely, about every egg rayed developed an abnormality which conformed to this general, U-shaped type. Three type-experiments may be presented for the purpose of demonstrating the constancy of results. On May 27, 1914, 9 eggs in the two-celled stage were rayed, each for 1 minute. One died before development had progressed more than 24 hours. Of the 8 remaining all were U-shaped. Three of these were permitted to develop until the tail and head were well formed. The others were killed at earlier developmental stages. On June 18, 1914, 25 eggs were rayed, each for 30 seconds. These were in the one- and two-celled stage. Various portions of the prescribed area above the equator were influenced. Three died within 24 hours, one lived until the neural groove had disappeared and presented a marked curvature of the trunk. All of the remaining 21 were U-shaped. On June 23, 1914, an exposure of 19 eggs in the one- and two-celled stage was made, each for 30 seconds. Eight of these were rayed in the designated area, and 11 along the equator. One of the eight died early, of the remaining 7 all were U-shaped. The 11 others demonstrated spina bifida.

The inference seems justifiable, therefore, that the constant type of defect produced is the result of a constant and uniform degree of alteration of the superficially-placed egg substance in the area illuminated. Just what the nature of this effect is we are at present unable to state. It would appear that the protoplasm, using this term in its broadest sense, had been modified in some chemical way to a degree which not only rendered it unfit for participation in the subsequent chemical ontogenetic

processes of which it normally was a part but, in addition, caused its presence to act as a mechanical hindrance to the developmental shifting of anlagen.

Studies of the action of ultra-violet light rays have produced proof that proteins, carbohydrates, and lipins may be chemically altered by means of this form of energy. To review the experimental results briefly: Massol and Kluver have ascertained that starches may be altered, and Stoklasa, Zdobnický, Chau-chard and Mazoné, and Pougnet that they may be ultimately broken up. Diastase may be liberated in plants (Maquenne and Demoussy). Bierry, Henri, and Ranc have inverted cane sugar. Agulhon, Maurain and Warcollier, and Raybaud presented evidence that the action of enzymes in the presence of oxygen was weakened and destroyed by ultra-violet light. Furthermore, albumen may be coagulated (Bovie) and the iodine content of fats may be lowered (Roemer and Sames). Thus we may reasonably infer that ultra-violet light rays are capable of bringing about certain chemical modifications of the protoplasm of the cell to all or to part of which we may attribute the abnormality in development noted. The evidence is lacking at present, however, which might enable us to associate this developmental result with a specific alteration of protein, of carbohydrate, or of lipin.

It was pointed out in the preceding paper that the superficial mass of altered protoplasm situated between the approximating neural ridges prevented their normal fusion with the consequent formation of a complete neural tube. The condition of spina bifida resulted, therefore, as a direct result of the mere mechanical interruption of this process. The action of the altered protoplasm was owing entirely to its passive function as a mass which mechanically interfered with the medianward, migratory movement of the hemineural anlagen. At least, in this one factor could be found sufficient evidence as the causative condition of this abnormality.

In the present instance of the U-shaped embryo, however, the inert mass was located lateral to and above the level of the hemineural anlagen. It could not, therefore, by virtue of interposition mechanically interfere with the approximation of the

neural tube-halves. Because of its lateral position and close relationship to one of these halves it might exert, however, a restraining influence upon the rate of migration of the tube-half of the corresponding side of the egg. This retardation of the shifting of the hemineural anlagen on the affected side induced a correspondingly exaggerated migratory process in the anlagen upon the unaffected side of the egg. As a result of this loss of coördinated, migratory movement, the anlagen upon the affected side had not yet reached the median plane before fusion took place with the unaffected anlagen by reason of their exaggerated migration. The normal angularity of the hemineural long-axis persisted more or less completely, therefore, in the adult tadpole as a result of this process. This was expressed by the curvature of the trunk invariably towards the affected side. The location of this flexure, as was stated above, was directly dependent upon the segmental level of the affected area of protoplasm.

It is a singular fact that the histological sections failed to demonstrate a single gross or microscopic defect in these embryos. The viscera developed normal in shape and in size. No defects could be recognized in the muscle segments. The mass of these in the concavity of the flexure was estimated roughly to be the same as that of the corresponding segments on the convex side of the body though their shape was necessarily distorted to fit the concavity. The yolk mass of the trunk of the embryo was completely utilized in the differentiation of the tissues and in the metabolic processes of the embryo. There was no evidence at any time that a portion of this yolk mass had been rendered completely inert by the rays and consequently excluded from participation in the normal chemical developmental processes of the body.

The action of the rays inferentially, therefore, had not induced such chemical changes as would permanently eliminate the substances affected from participation in the normal organo-genetic or somatogenetic processes. But the rate of chemical modification of these substances in participation in the normal metabolic processes of the embryo was, however, retarded. We may find in this retardation of chemical participation additional evidence for the causation of the check of the normal lengthening process

of the embryo on the affected side. The opposite side progressed approximately at its normal, but comparatively greater, rate. As a result, the flexure of the neural tube increased proportionate to the developmental progress of the embryo. In other words, as an embryo increased in age the angle formed by the trunk with the tail became more and more acute until a condition was arrived at finally in which the long axis of the trunk lay parallel to that of the tail.

The sketches presented with this paper are intended merely to demonstrate the general morphological and histological features which are characteristic of the development of these abnormal, U-shaped embryos. It may be stated, however, that no great attempt has been made to make them faithfully representative of all of the histological details of the tadpoles. In many instances the drawings are composite. In every instance, however, the basis of each drawing has been an actual specimen. The figures are to be interpreted, therefore, as merely type representatives of the numerous examples which are illustrative of the peculiarities of this form of abnormal development.

Fig. 2 represents an embryo in the early stages of development




FIG. 2.

viewed from the dorsal aspect. The deep furrow which penetrates from the left, to and beyond the center of the embryo mass represents the line of apposition of the two portions of the right flank of the embryo which is so bent upon itself that the right surface of the tail region is in contact with the right surface

of the head region. Actual continuity of the tissues, however, does not exist. The cephalic end of the embryo, which is represented below and to the left of the sketch, is distinguished by its greater transverse diameter. The area of protoplasm in this specimen affected by the rays lay in the concave bend of the trunk at the extremity of the cleft. It will be noted that this flexure is sharply restricted to a relatively small portion of the embryo. Cephalic to and caudal to this level, both the head-end and the tail-end of the embryo are, respectively, of normal shape.

Occasionally, the two neural tube-halves fail to unite in the caudal region, bringing about a condition which is demonstrated well by Fig. 3. In this the yolk mass is represented as projecting




FIG. 3.

to the right between the caudal extremities of the divided neural tube. This is an early developmental feature in the production of the condition of spina bifida, the divided condition of the neural tube-halves persisting in the adult free-swimming tadpole. The bend in this particular embryo is represented as occurring high up in the trunk region. The cephalic end of the embryo is shown with its right surface resting against the right neural tube-half caudally. As was noted in connection with the preceding figure, so is it likewise true of this, both cephalic to and caudal to the angle of the fold, aside from the features of spina bifida, the appearance of the embryo is practically normal. The stages of development represented by these two figures are relatively early, consequently but little of the ordinary features of external configuration of the embryo can be made out.

A still later stage of development demonstrating a more marked differentiation of the tail from the head region is shown by Fig. 4, while the tadpole in Fig. 5 represents the external features of these tadpoles during the free-swimming period. In this latter picture, the sucker, the eyes, the external gills and the




FIG. 4.

tail fins are readily made out. The sharpness of the fold in the trunk in both instances indicates a restricted localization of the alteration produced in the protoplasm of the developing ovum.

These creatures are unable to straighten out the flexure of the trunk. They swim, as was noted before, by means of ordinary tail movements, the vigor of which is not impaired




FIG. 5.

in any single instance. As can readily be understood by reference to the figure, however, the abnormal position of the caudal long-axis, with regard to the long-axis of the head and remaining portion of the trunk, is such that the swimming movements of the tail tend to force the embryo about in a circular or spiral

direction. Because of the condition of fixity of the bend progression directly forward is necessarily out of the question.

The deformity represented by Fig. 6 is illustrative of the second mechanism of production of the folded embryos. The concave side of the neural tube, to the right in the figure, is occupied by irregular bulgings of the epidermal surface of the embryo. When the development of embryos conforming to this type is followed from day to day, it is noted that the flexure of the neural tube becomes greater from day to day. The apparent occasion



FIG. 6.

of this is referable chiefly to two factors; one, the normal but relatively more rapid growth of the normal myomeres to the left of the longitudinal axis of the body, and the other, to the relatively, and actually slower, growth of them upon the concave side of the median body plane. Histological studies demonstrated as well that differentiation in the myomeres upon the affected side was somewhat retarded.

There are, therefore, these two general developmental mechanisms either of which leads to the production of a U-shaped embryo. In one the angularity of the bend is constant from the time of its first appearance in the embryo, whereas in the other this angularity increases with the succeeding stages of differentiation and growth of the anlagen. The end result is identical, however, in both instances both from the morphological and the histological standpoint. Furthermore, in both instances the segmental relationship of the bend to the area of protoplasm altered in the ovum is the same.

The broken lines shown in Fig. 4 bear numerals which correspond to the planes of section represented by the succeeding figures, 7 and 8. The cross-section represented by 7, therefore, passes through the embryo at the level of the fold in the neural tube. The two halves of the yolk mass with sections of the




FIG. 7.

enteron which belong, the one to the cephalic end and the other to the caudal end of the embryo, are shown separated from each other by a deep cleft lined by ectoderm. Dorsal to the enteron lie two sections of the notochord with the myomeres sectioned in their long-axis between them. Between and dorsal to these




FIG. 8.

two portions of the notochord, the oblique section of the bend in the neural tube, demonstrating the neurocele as well, is to be seen. The level represented by Fig. 8 includes the region of the brain and subjacent pharynx to the right and the trunk to the

left. Both portions of the embryo are connected with each other by a narrow strip of mesenchymal tissue at the bottom of the epidermal cleft. The notochord is to be seen only in the left half of the embryo with the myoblastic tissue to either side of it, the enteron ventral and the neural tube dorsal. The stage is so early that most of the yolk cells are in an undifferentiated condition.

Figs. 9, 10 and 11 are sections of one and the same embryo.




FIG. 9.

In this the bend of the trunk occurred at about the level of its middle segment. The section-knife encountered both the neural tube and the notochord at their curvature. The undifferentiated yolk-mass of the two limbs of the U are connected by a relatively slender commissure-like mass of yolk cells and of



FIG. 10.

mesenchymal tissue. The section of Fig. 10 taken at about the level 8 of Fig. 4 demonstrates on the left the transsected notochord with the cephalic end of the definitive spinal cord dorsal to it, both imbedded in myoblastic cells not yet well differentiated. A section of notochord and of spinal cord with the

enteron ventrally placed to them are to be seen in the right half of the sketch. There is no indication at this early stage of a beginning formation of the anlagen either of the liver or of the pancreas. The section represented by 11, however, passes through the head region at the level of the eyes. To the left in this sketch there can be identified the cephalic portion of the neural tube with its well-differentiated walls showing a degree of stratification. The neurocele appears clear and contains no cell




FIG. 11.

detritus. The neuroblasts do not demonstrate any pathological features. A thin layer of mesenchymal tissue intervenes between the roof of the neurocele and the epidermis. Immediately ventral to this section of the brain cavity the knife has encountered another portion of the normally folded brain stem with its ventricle and, projecting to either side from its ventral aspect, a portion of the optic commissure and stalks proceeding laterally to the optic cups. The cytological appearance of these structures presents no departures from normal. The cavities are free from cell detritus. The lens of the eye on both sides is in its normal position. Stratification of the walls of the optic cup, of the stalk and of the brain vesicle have progressed to their normal relative degree. Immediately ventral to these structures in the median line is shown a section through the pharynx, the cells of which, as well, show none but normal features. The right half of the sketch demonstrates the prominent caudal fin, transected neural tube and notochord, and the caudal extremity of the enteron, the last embedded in the dorsal region of the yolk

mass. The mesenchymal tissue to either side of the neural tube and notochord presents normal features.

Comparable to the levels represented by Figs. 7 and 9 is the section represented by Fig. 12. Here again the flexure of the neural tube lies dorsal to the notochord sections. Immediately




FIG. 12.

ventral to the latter the forms of the elongated myoblasts are to be made out. A relatively early stage of development of the enteron is evidenced by the condition of the yolk cells, the symmetrical masses of which are separated by the deep epidermal cleft.

In the succeeding four figures, 13, 14, 15 and 16, embryos are




FIG. 13.




FIG. 14.

represented in each of which the angularity of the flexure was not sufficient to bring the long axis of the two body portions parallel to each other. In each instance, the cephalic extremity

of the embryo projected from the trunk at approximately a right angle. Consequently the neural tube is sectioned in the head region parallel to its long axis, but in the trunk region at right angles to its long axis. In Fig. 14, for instance, the head end of the embryo projects to the right. The walls of the brain




FIG. 15.

vesicle demonstrate no abnormal features. Because of a certain amount of twisting of this extremity but little of the pharynx appears in the section upon the ventral aspect of the brain stem. At the left extremity of the neural tube an almost perfect tran-



FIG. 16.

section of the cephalic end of the cord is shown. The notochord with the transected enteron and the yolk-cell mass are represented immediately ventral to the tube. In Fig. 15, the cephalic

extremity of the embryo is shown above to the left resting upon the trunk of the embryo. Here again approximately the long axis of the brain tube has been encountered by the section knife. Stratification of brain walls has begun. The enteron together with the notochord lie to the right in the sketch. The mesodermal cells constituting the wall of the pharynx lie immediately ventral to the brain tube. Fig. 16 presents these same general features and, in addition, a bifid appearance of the caudal portion of the neural tube upon the right side of the sketch. The ectoderm folds in between the two tube-halves each of which, while presenting an entire tube, demonstrates its derivation by reason of the thinning of its medial wall. This appearance has been previously referred to in the paper on the condition of spina bifida.

It is to be noted that not one of the histological sections, which are presented, as was mentioned before, as merely typical instances of the appearance of the tissues during the different developmental stages of the embryos, presents any indications whatever either of cell or of tissue disorganization. There is no indication of the presence of degenerated areas of protoplasm either in the yolk mass or in the normal body cavities, nor can there be found any degenerated or extruded nuclei. The organs of the embryo such as the enteron, the brain, the cord, the myomeres, the notochord, and so on, seems to be normally formed and are of the normal size. So far as can be ascertained through the study of longitudinal sections of the myomeres at the level of the bend in the trunk the cause of this phenomenon is not, apparently, referable either to a degeneration or to a falling out of myomeres upon the concave side of the trunk. As might be reasonably expected, however, the myomeres upon this side have a form which departs from the normal in so far as the segments conform to the concavity. The corresponding segments on the opposite side of the fold are consequently relatively elongated. The normal morphological appearances of the tissues argued for a normal physiological activity as well, and this seemed to be attested by the movements and general vitality of the tadpoles. None of these tadpoles was permitted to live, however, to the period of transformation.

In the preceding paper on "The Artificial Production of Spina Bifida" it was noted that, apparently, the proanlagen and anlagen of the embryo were restricted to the pigmented hemisphere of the egg, that the raying of a small localized surface-area of the yolk hemisphere or of the region of the quarter produced invariably the condition of spina bifida in the embryo. It was noted in these embryos, however, that indications of the destruction of cell groups or of definite areas of protoplasm in the adult tadpoles were absent. No gross structural defects of organs aside from the bifid character of the spinal cord were demonstrable. Where the dosage of ultra-violet light ray energy was sufficiently great, however, a mass of protoplasm equal in size to the superficial area affected was extruded from the body of the embryo. This was observed as well in this present experiment. Under such conditions naturally the developed embryo demonstrated both an organic and a structural defect. When the dosage is reduced in amount, however, it is demonstrated both by these and by the spina bifida experiments that the possibility exists for the production of a developmental defect the structural cause of which may not be present or recognizable in the adult. The absence of atypical cytological characters as well as organic defects and of exovates must force us to the conclusion that either such did exist for a longer or shorter period during the development of the embryo and subsequently became incorporated into the structure of the body as the result of an elaborate but delayed chemical transformation, or that there ensued upon the raying a transitory suspension of the physiological activity of the cells affected with a later complete resumption of that activity. Either conclusion must force us to presume, however, the presence of a chemical alteration either of cell content or of cell mass as a direct result of the raying. Indeed, it would be rather difficult to contend that no change was brought about in the chemical composition of the protoplasm in the area affected by the rays.

Since it has been definitely proved that protein substances, carbohydrates and lipins may be altered to some degree in their chemical constitution by this form of energy, it does not appear to be unreasonable to infer that one or all of the corresponding

groups of substances may be susceptible to the influence of the rays while still in the living ovum. Exovation as the direct result of the application of a great amount of energy may be cited as an instance of this alteration. The normal histological features of the embryo, and of the adult tadpoles, however, argue the presence of a process of restoration or of regeneration during the time of development of the embryo of the changes which were brought about through the activity of the rays. It might be assumed, accordingly, that the chemical modification thus artificially produced is gradually rectified during the developmental cycle but during the early stages of this process the substances involved are incapable of entering into their normal functions in the chemical elaboration of the pro-anlagen and anlagen. The resultant action of these altered substances might be interpreted, therefore, as that of an inert body which serves, as was seen to be the case with the spina bifida embryos, as a check to the gross, mechanical shiftings characteristic of the normal genesis of the anlagen. The absence of an area of altered protoplasm or of organic defects in the adult tadpole is the basis for the logical argument that the chemical readjustment of the protoplasm, altered by the rays, is ultimately completed to the degree that this protoplasmic mass does enter finally into its usual function in the elaboration of the normal anlagen.

BIBLIOGRAPHY.

Agulhon

'12 Ann. Inst. Pasteur, XXVI., 1, p. 38.

Baldwin

'15 Anat. Rec., IX., 5, p. 365.

Bierry, Henri, and Ranc

'11 Compt. rend. Acad. Sci., (Par.) CLII., 23, p. 1629.

Bovie

'13 Science, XXIV., p. 373.

Cercelet

'10 Rev. vit., XXXIII., 842, p. 124.

Chauchard and Mazone

'11 Compt. rend. Acad. Sci., (Par.), CLII., 24, p. 1709.

Euler

'12 Abst. in Ztschr. Bot., IV., 6, p. 450.

Henri

'12 Compt. rend. Acad. Sci., (Par.), CLV., 4, p. 315.

'14 Compt. rend. Acad. Sci., (Par.), CLVIII., 14, p. 1032.

'14 Compt. rend. Acad. Sci., (Par.), CLIX., 4, p. 340.

Henri, Henbronner, and von Recklinghausen

'12 Abst. in J. Soc. Chem. Indus., (Par.), XXXI., 2, p. 1004.

Henri and Moycho

'14 Compt. rend. Acad. Sc., (Par.), CLVIII., p. 1509.

Kluyver

'11 Sitzber. K. Akad. Wiss. (Vienna), Math. Naturw. Kl., CXX., 1, no. 10,
p. 1137.

Maquenne and Demoussy

'09 Compt. rend. Acad. Sci., (Par.), CXLIX., 19, p. 756, and 22, p. 957.

Massol

'11 Compt. rend. Acad. Sci., (Par.), CLII., p. 902.

'11 Compt. rend. Soc. Biol., (Par.), LXX., 13, p. 509.

Maurain and Warcollier

'09 Compt. rend. Acad. Sci., (Par.), XLIX., 2, p. 155.

Pougnat

'10 Compt. rend. Acad. Sci., (Par.), CLI., 12, p. 566.

Raybaud

'10 Compt. rend. Soc. Biol., (Par.), LXVIII., 15, p. 772.

Roemer and Sames

'10 Hyg. Rundschau, XX., 16, p. 873.

Stoklasa

'11 Centbl. Bakt., XXXI., 2, p. 477.

'13 Abs. in Bot. Centbl., CXXII., 4, p. 90.

'14 Ztschr. Pflanzenkrank, XXIV., 4, p. 193.

Stoklasa, Sebor, Zdobnický

'12 Biochem. Ztschr., XLI., 5, p. 333.

Stoklasa and Zdobnický

'10 Chem. Ztg., XXXIV., 107, p. 945.

'11 Bioch. Ztschr., XXX., 6, p. 433.

DIFFERENTIAL SUSCEPTIBILITY AS A BASIS FOR MODIFICATION AND CONTROL OF EARLY DEVELOPMENT IN THE FROG.

A. W. BELLAMY.

CONTENTS.

I.	Introduction and General Statement of the Problem.....	312
II.	Materials and Methods.....	314
	1. The Susceptibility Method.....	314
	2. Routine and Methods.....	317
III.	The Origin of Polarity in the Egg.....	318
	1. Polarity in Other Organisms.....	318
	2. Origin of Polarity in the Frog's Egg.....	319
IV.	Experimental Data.....	322
	1. Susceptibility to Lethal Concentrations of External Agents.....	322
	(a) Unsegmented Eggs.....	323
	(b) Cleavage Stages.....	324
	(c) Gastrula Stages.....	325
	(d) Later Embryonic Stages.....	326
	2. Experimental Modification and Control of Development.....	328
	A. Disturbances in Cleavage Such That the Egg Never Gastru- lates.....	332
	(a) Cell Size Ratios.....	332
	(b) Bilaterality.....	333
	(c) The Segmentation Cavity.....	333
	(d) Meroblastic Cleavage.....	335
	B. Disturbances in Gastrulation.....	335
	(a) Retardation of the Dorsal Lip Region.....	336
	(b) and (c) Retardation of Both Dorsal and Lateral Lip Regions and Equatorial Gastrulation.....	337
	(d) Secondary Invaginations.....	341
	(e) The Gastrular Angle.....	343
	C. Differential Inhibition in Later Development.....	344
V.	Discussion.....	346
	1. The Nature of Physiological Axes.....	346
	2. Origin of the Dorsal Lip Region.....	349
	3. The Question of "Specificity" in Vertebrate Teratogeny.....	351
VI.	Summary.....	356
VII.	Literature.....	358

I. INTRODUCTION AND GENERAL STATEMENT OF THE PROBLEM.

Observations on vertebrate teratology date almost from the beginning of descriptive biology, though active accumulation of experimental data on the frog especially, began little more than

a quarter of a century ago. Among the early papers on the subject may be mentioned those of O. Hertwig (1892, 1894, 1895, 1896); Gurwitsch (1895, 1896); Morgan and Tsuda (1894).

These and other workers have studied the early development of certain teleosts, amphibia and birds under the influence of such physical factors as: gravity, mechanical pressure, temperature, light, electricity and magnetism, X-rays, radium emanations, atmospheric pressure, and a wide variety of chemical substances.

It is perhaps sufficient, here, merely to mention the essential similarity of terata produced by widely different methods, such, *e.g.*, as high and low temperatures, many different chemical agents, and heterogeneous hybridization. Equatorial gastrulation, embryos with permanent yolk plugs, with spina bifida, microcephalic forms and many other abnormalities in the frog have been produced by nearly all the methods known to experimental teratology. For instance, microcephalic frog embryos with permanent yolk plugs, and spina bifida have been produced by exposing different stages in development to such external agents as: high temperature (Jenkinson, 1909, p. 98, fig. 49); 0.6 per cent. NaCl (Hertwig, 1894, p. 315, 316, 1895, plate XX., figs. 5, 6, 7, 10, 12, 15, 16, et al); fertilizing "over-ripe" eggs (Hertwig, 1892); strychnine (Gurwitsch, 1896); sugar solutions (Bataillon, 1901, figs. 2, 5); Jenkinson (1906) produced these same types (among many others) in a number of isotonic solutions of Na_2SO_4 , K_2SO_4 , NaNO_3 , urea, sugar solutions, et al.

Cyclopia in *Fundulus* has been produced by physical and chemical agents and by hybridization. Stockard (1909) obtained this anomaly with the aid of magnesium chloride and for a time ascribed a specific cyclopia producing property to this salt. Later, Stockard produced cyclopic monsters with alcohol and other substances and was forced to abandon his theory of specificity. McClendon (1912) had no difficulty in producing cyclopia in isotonic solutions of NaCl, LiCl, MgCl_2 , NaOH and in alcohol, and other substances. McClendon is somewhat at a loss to explain his results but sums up the facts in these words: "We need only suppose that the cells between the eye anlagen are more easily affected than other cells of the embryo, to explain

the action of the solutions." Werber (1915, 1916, 1917) obtained all the known types of terata in *Fundulus* by using substances usually found among the products of destructive metabolism, viz., butyric acid, acetone, urea, lactic acid, etc. In the concentrations used he found butyric acid and acetone, especially the latter, somewhat more effective than the other substances in the production of monstrous development. Werber looks upon abnormal development as due to a "blastolytic" action of the deleterious agent upon the primordial regions of the eggs, especially those at the anterior end of the body. Kellicott (1916) has secured the usual wide variety of terata in *Fundulus* simply by subjecting the early egg stages to the action of low temperature. He is inclined to look upon teratogenesis as being due in some way to a "disorganization" of the egg materials. Newman (1917) noticed the same range of abnormalities among his heterogeneous *Fundulus* hybrids and applied Child's axial gradient theory to the interpretation of these and other terata seen in teleosts produced by chemical and physical agents.

Perhaps enough has been said by way of a preliminary statement to indicate the essential similarity of experimentally produced monsters regardless of the methods employed in obtaining them. Likewise a little reflection on the various theories advanced to account for monstrous development makes apparent the lack, hitherto, of an adequate basis for a rational interpretation of them.

It is the purpose of this report to attempt an analysis of the problem of physiological axiation and teratogeny in the frog from a physiological point of view. It considers the general problem of physiological axiation, the origin of polarity, and presents data on the modification and control of early development in the frog on the basis of differential susceptibility.

II. MATERIAL AND METHODS.

1. *The Susceptibility Method.*—Observations on many species of animals and plants have shown that definite and characteristic differences in susceptibility to a wide variety of physical and chemical agents, within certain limits of concentrations or in-

tensities of action, exist in relation to the physiological axes.¹ These differences in susceptibility may be demonstrated in several ways: first, as differences in survival time of one region of the egg or embryo as compared to other regions, under conditions severe enough to kill without permitting acclimation to occur; second, as differences in the degree of inhibition of growth and development, or in certain cases as differences in the degree of acceleration of these processes; third, as differences in the rate or degree of acclimation to a certain range of less severe conditions; fourth, as differences in the rate or degree of recovery after temporary exposure to conditions that inhibit development.

These differences in susceptibility determined in these different ways are all expressions of the fact that a "differential susceptibility" to the action of external agents is a characteristic feature of physiological axes in both plants and animals so far as they have been examined with reference to this point. This "differential susceptibility" appears according to experimental conditions as differential disintegration associated with death, and as differential inhibition, acclimation or recovery in development.

The high degree of uniformity in the susceptibility relations in different organisms, both animals and plants, to a wide variety of agents and conditions, viz., cyanides, anesthetics, acids, alkalis, various salts, certain alkaloids, physical conditions such as extremes of temperature and certain conditions that may be termed negative, such as lack of oxygen, indicate that in their general features these susceptibility relations are independent of specific qualitative differences in the protoplasm of different forms. If this is the case they must depend upon quantitative differences of some sort which are common at least to all forms in which these differences in susceptibility have been shown to exist.

So far as present knowledge goes, the facts indicate that susceptibility, in the sense in which the term is used here, is

¹ The extensive literature on this subject, previous to 1915, mostly the work of Child, is treated in two of his recent books (1915*a*, 1915*b*) to which the reader is referred for specific references. Other references are: Behre (1918); Child (1916*a*, 1916*b*, 1916*d*; 1917*a*, 1917*b*, 1917*c*, 1917*d*; 1919*a*, 1919*b*); Child and Hyman (1919); Hyman (1916*a*, 1916*b*; 1917*a*, 1917*b*; 1919).

associated in some way, directly or indirectly, with the rate of certain fundamental metabolic reactions, such, *e.g.*, as oxidation-reduction processes, and the protoplasmic conditions associated with them. Considerable evidence has been accumulated which indicates that in general the susceptibility to lethal or strongly toxic conditions and the capacity for acclimation to less severe conditions and for recovery after temporary exposure—all vary more or less directly with, though not necessarily proportionally to, the rate of fundamental metabolic reactions. This does not mean that all the agents employed act directly upon these reactions in every case or that all act in the same way upon protoplasm. It probably means merely that living protoplasm is a system of more or less closely associated and interdependent reactions and conditions so that no essential factor in this system can be altered beyond a certain degree without involving the system as a whole.

But, however the facts of differential susceptibility may finally be interpreted, concerning the facts themselves there can be no doubt. They have afforded a means not only of demonstrating characteristic differences in physiological conditions along the axes of organisms but also of modifying and controlling development in definite ways through these differences. This study of the development of the frog was undertaken in the attempt to determine whether, and to what extent, the susceptibility method could be used in modifying and controlling development in a vertebrate, as they have been used in investigations of the physiology of development in certain invertebrates.

I am under obligation to Professor C. M. Child, in whose laboratory this work was done, for essential aid in the way of suggestion and criticism, and to Dr. L. H. Hyman for many thoughtful suggestions. Work on amphibia involving the susceptibility method was begun by Dr. Child in 1913. In the spring of 1916, work was begun by him on the modification and control of development in the frog, on the basis of differential susceptibility; the data and preserved material of these experiments were turned over to me late in that year. My own work was done in the spring of 1917 and the spring of 1919 during which time approximately one hundred thousand eggs were handled, of which about forty thousand were preserved.

2. *Routine and Methods of Handling Eggs.*—Clasping frogs (*Rana pipiens*) were obtained from a local dealer and placed in a large cement aquarium arranged to simulate natural conditions as nearly as possible. The temperature of the water was kept between 5° and 9° C. There was no difficulty in obtaining all the eggs that could be used; and with a little patience they could be had as soon after deposition as desired. As many as five females have been seen spawning at one time. Shortly after deposition each egg mass was given a number, the jelly cut into bits containing ten to twenty eggs each, and then placed in a gallon jar of the cold aquarium water. About an hour before being introduced into experimental conditions the jar of eggs was removed to the laboratory to warm slowly up to the room temperature ($17 \pm 1^\circ$ C.).

For the modification of development the following chemicals were used: potassium cyanide, formaldehyde, potassium permanganate, mercuric chloride, magnesium chloride, lithium chloride, hydrochloric acid, sodium hydrate, and ethyl alcohol. The method of handling the eggs varied somewhat with the chemical used. Eggs treated with KCN, CH₂O, HCl, NaOH and C₂H₅OH, were placed in liter Erlenmeyer flasks filled almost full and stoppered. For the other chemicals, glass finger bowls of 300-c.c. capacity, one- and two-liter candy jars were used and covered with glass plates to prevent evaporation.

In a given experimental series similar glassware was used, each vessel containing approximately the same number of eggs from the same female. In every case one vessel of the series was used as a control and was treated in the same way as regards stoppering, changing of fluids, etc. Eggs from the same females developing in liter Erlenmeyer flasks filled nearly full and stoppered, and in the same volume of water in open dishes did not differ in rate or manner of development up to some time after hatching, provided the water was changed once in twenty-four hours and when there were not more than 100 c.c. of egg mass in the stoppered flask. Especial care was taken to use no more than 15 to 20 c.c. of egg mass in stoppered vessels, and to change solutions in all experiments daily.

As will be noted later, the frog egg undergoes a tremendous

increase in susceptibility during the early stages of development, and to provide for this and to carry the effects of differential inhibition somewhat farther than is possible where the eggs remain in the same concentration throughout the experiment, a number of experiments were done where the solution was gradually diluted as development proceeded.

In the experiments on recovery, the eggs were removed from the solutions and washed in several changes of water to remove any of the chemical remaining in the jelly.

III. THE ORIGIN OF POLARITY.

I. *Polarity in Other Organisms.*—There is abundant evidence to show that for many organisms and especially among the plants, polarity and symmetry arise in response to external conditions. Polarity in the egg of the alga *Fucus*, and in the spore of *Equisetum* is usually determined by the direction of incident light. In some algæ, polarity and symmetry are directly under the influence of light, even in the vegetative thalli, and are reversible or modifiable by change in light relations. In various liverworts and free prothallia, and even in certain phanerogams, light is an important factor in determining dorso-ventrality in branches.

The polarity of a number of plant and animal eggs bears a definite relation to their manner of attachment to the parent body during growth stages. In the phanerogams, for example, the free end of the egg becomes the apical, the attached end the basal end of the plant axis, and the same relation holds for a number of the lower animals.

Likewise a polar axis of symmetry once established may be obliterated experimentally and a new one induced by conditions external to the organism. This has been done by H. V. Wilson (1907, 1911) in certain sponges and hydroids, work which was confirmed by Hargitt (1915). In experimentally produced bi-axial forms in planaria, hydroids and annelids, a new axis arises 180° from the old one. In *Corymorpha*, e.g., Child showed that when pieces of the stem are placed in 2 per cent. to $2\frac{1}{2}$ per cent. alcohol in sea water, "in the course of a few days the pieces become shorter and more rounded, decrease in size, and lose the

characteristic structure of the *Corymorpha* stem." On removal to water after several days in alcohol, a new individual arises from the old tissue with its axis at right angles to the old one.

Various investigators have observed that under certain conditions, *Hydra* may lose its characteristic structure and "melt" down into a shapeless mass from which a new individual may arise. Dr. Hyman, in this laboratory, has recently observed that such masses often give rise to several hypostome regions each with tentacles.

Physiological axiation begins, of course, with the origin of polarity in the egg. Polarity may, conceivably, arise in several ways. First, axiation and polarity may, as has been commonly supposed, exist in protoplasm in relation to some intrinsic molecular or other structure which is a fundamental property of living matter; second, the primordial cell, from which the ovum arises, may inherit its polarity, which, if true, merely pushes the problem farther back; third, the appearance and position of local differences in structural and functional order in the egg protoplasm, that constitutes polarity, may arise during the growth and development of the egg in response to conditions external to it. The first alternative has been shown to be untenable, and the second offers so little satisfaction, that, in the light of the accumulated evidence, it seems necessary to consider the last alternative as the more probable one, at least until it has been shown that polarity arises in some other way.

2. *Origin of Polarity in the Frog's Egg*.—The mature egg of the frog is described as possessing a radial symmetry about an imaginary axis, the polar axis, passing approximately through the centers of the pigmented (animal, apical) hemisphere and of the unpigmented (vegetative, basal) hemisphere. This polarity is marked by the localization of most of the protoplasm in the pigmented hemisphere, by the eccentricity of the nucleus, and by the distribution of pigment. In order to determine, if possible, whether, in the frog egg, a relation exists between its polarity and conditions external to it, a study was made of the ovarian relations of the egg, especially of the relation of the polar axis to the region of attachment to the ovarian membrane and to the blood supply of the egg.

Some days or weeks before extrusion, the egg is suspended from the ovarian membrane in an epithelial sack (Fig. 1). This sack, the theca, forms at one region on the egg a narrow stalk, the pedicle, which becomes continuous with the ovarian membrane. Beneath the theca is a layer of follicular cells and beneath this, the vitelline membrane. In the theca are found the blood vessels that supply the egg.

Table I. gives the location of the pedicle with reference to

TABLE I.

Location of Pedicle.	Number of Eggs Observed.	Per Cent.
Position A (see Fig. 2).....	75	12.2
Position B.....	217	35.3
Position C.....	259	42.2
Position D.....	63	10.3

These results were obtained by taking a number of eggs at random from various parts of the ovary and determining the location of the pedicle for every egg in each sample.

the distribution of the pigment on the egg. It will be noted that in approximately eighty per cent. of the cases, the region of attachment (the pedicle) is near or on the boundary between the pigmented and unpigmented regions of the egg (Fig. 2).

A number of specimens were injected through the conus arteriosus with a lead chromate-starch-gelatin mass¹ and in the great majority of cases only those vessels lying over the unpigmented hemisphere remained uninjected and could often be traced by the color of the blood remaining in them. In several cases injected vessels were seen to pass a little distance over the unpigmented hemisphere, but in no case was an egg observed where more than a small per cent. of the vessels overlying the yolk were injected (Fig. 3). Several unsuccessful attempts were made to inject the venous system.

In order to eliminate any error due to the injection mass passing through the capillaries into the veins, several frogs were opened under normal salt solution and the circulation of the blood over the egg determined with the aid of the binocular microscope. In every case where the movement of corpuscles over the pigmented hemisphere could be seen, the circulation

¹ Guyer (2d edition, page 84). The mass was made up in a 1 per cent. uncooked starch suspension, warmed and well stirred immediately before the injection was made.

was away from the heart and therefore arterial. The movement of blood over the unpigmented hemisphere, where the movement of corpuscles could be seen more clearly than over the pigmented hemisphere, was always toward the heart and therefore venous. As was noted for the injected specimens, a small per cent. of the eggs showed arteries passing a little distance over the yolk. But it is especially to be noted that in every case observed, the greater part of the arterial blood supply was restricted to the pigmented hemisphere.

In the ovary of the frog, the polar axes of the eggs lie in every possible direction with respect to gravity, so that the possibility of polarity being determined in the ovary with respect to gravity is at once eliminated.

The data indicate that polarity in the frog egg arises at some time during the growth stages, in response to external conditions, viz., to the blood supply of the egg: that region of the oögonium chancing to be most richly supplied with arterial blood being destined to become, by virtue of this respiratory and nutritive relation, the animal pole of the egg. Naturally, the region on the surface of the egg where the capillary net work is most extensive would be effective in determining polarity rather than the point where the blood vessels enter the theca.

A region of higher oxidation rate in the egg, or at any rate the proximity of this region to a greater oxygen supply (arterial blood) is further evidenced by the appearance of pigment—a melanin (Kellicott)—over a limited surface of the egg, for it is well known that oxygen is necessary for the formation of these pigments. It is probable that the origin of polarity and the appearance of pigment in a symmetrical relation to that polarity are both expressions of the localized oxygen supply (or nutritive supply, or both) of the egg.

It may be noted here also that pigment appears most densely in the most active regions of the egg where other evidence indicates that oxidations are proceeding more rapidly than elsewhere. In other words, the density of pigmentation seems to be an expression of the rate of at least certain oxidations occurring in that region. Local increase in activity results in the formation or increase of the pigment and local decrease in activity results in diminution of pigment. Those unpigmented cells lining the

archenteron in the early gastrula stages, *e.g.*, become pigmented and most densely where cell division is most rapid.¹ And to anticipate somewhat, the subjection of early gastrula stages to strongly inhibiting conditions, results in a marked diminution of pigment in the dorsal lip regions.

The data further indicate that the polar axis as well as other axes that arise later in development are primarily gradients in fundamental physiological conditions or processes, which are




FIG. 1. Semi-diagrammatic view of a section through the germinal vesicle and pedicle of a full grown ovarian egg. *b.v.*, blood vessel in the theca; *f*, follicle cells; *g.v.*, germinal vesicle; *p*, pedicle; *th.*, theca.

made evident in several ways, such as, the differential cleavage rate which results in a gradient in cell size, by gradients in susceptibility, etc. (see pp. 322-328).

Discussion of the question of whether the location of the pedicle and the localized blood supply is related in any way to the definitive bilaterality remains for future consideration.

The fact that the pedicle is equatorial in position is of course, merely a necessary consequence of the way polarity and the accompanying distribution of pigment is determined by the blood supply.

IV. EXPERIMENTAL DATA.

1. *Susceptibility to Lethal Concentrations of External Agents.*

The early stages of the frog egg are extremely resistant to toxic agents. Eggs just beginning to segment, when placed

¹ King (1902).

e.g., in $m/10,000$ HgCl_2 , are accelerated in development for a short time, while gastrulating eggs die within a few hours (four to six) in $m/2,000,000$ HgCl_2 , without having developed farther after being placed in the solution. The same relation holds for other agents. Eggs from different females may differ consider-




FIG. 2. Diagrams of full grown ovarian eggs to show the position of the pedicle in relation to the distribution of pigment. (See also Table I.)

ably in their behavior when exposed to toxic agents; also eggs from the same female may differ somewhat in susceptibility to toxic agents but to a much less extent than in the former case. Some difficulty is experienced in watching the progress of disintegration in early stages, especially in concentrations that kill rapidly, on account of the spreading of the disintegrated protoplasm underneath the inner membrane, and thus obscuring the rest of the process. However, the progress of disintegration, wherever observed, is an orderly process and bears a definite relation to the polar axis of the egg and to the axes of symmetry of the embryo.

(a) *Unsegmented Eggs.*—The most satisfactory method found

for observing disintegration in the unsegmented egg, was to expose the egg soon after deposition, to a temperature of 0° C. for ten days or two weeks. Fig. 4 illustrates several different stages in the process of disintegration under such conditions. The eggs were deposited in the aquarium (temperature, 6.5° C.) March 26, 9:00–9:15 A.M., and were placed in the refrigerator at 10:00 A.M. On April 7, the eggs were in the condition shown in the figures. The relation of the disintegrated areas to the grey crescent and to the polar axis is obvious.

(b) *Cleavage Stages.*—The process of disintegration during early cleavage has been studied at low temperature, in KNC, $m/1,000$; $m/100$; HgCl_2 , $m/10,000$; in alcohol, 5 per cent., 8 per cent.:




FIG. 3. Semi-diagrammatic view of ovarian egg showing the distribution of arteries and veins. The blood vessels were outlined under the camera lucida.

and incidentally in all of the other agents used to modify development. The process of disintegration in all of these cases is essentially similar, except in the case of alcohol, which has a marked solvent action on the yolk. In any concentration of alcohol above 1 per cent., the yolk becomes rough and pitted, an appearance that is much accentuated in the higher concentrations (5 per cent. to 8 per cent.). Disintegration usually begins in a meridian that bisects the grey crescent and near the center of the pigmented hemisphere. The surface membranes of individual cells in this region break down and turn white. Shortly afterward, disintegration begins in the equatorial region just

above the middle of the grey crescent and spreads equatorially and apically to join the area where dissolution first began. At this time the disintegrated area forms, roughly, a triangular figure whose apex is the point where disintegration first began, and whose base is the equatorial region immediately above the grey crescent. Disintegration then spreads more or less symmetrically from the apex and sides of the triangle over the rest of the animal pole, and from the base to involve the grey crescent




FIG. 4. Unsegmented egg seen from above (apical pole) showing different stages in disintegration after twelve days' exposure to a temperature of 0° C. Disintegrated areas are represented by stippling. The dotted circle represents the boundary between pigmented and unpigmented hemispheres. The dotted crescent outlines the grey crescent. The cross marks the animal pole.

region. Fig. 6 illustrates the process just described. The history and treatment of the eggs is given in the legends to the figures.

(c) *Gastrula Stages*.—Eggs in an early gastrula stage, when placed in lethal concentrations of a toxic agent, always begin to disintegrate first in the dorsal lip region, and shortly afterward in the same meridian about 120° to 130° above the blastopore. From this upper point dissolution of the surface cells proceeds down the meridian and meets the disintegrated area of the dorsal

lip region, which has spread apically and now includes the lateral lips. The area then spreads more or less symmetrically from the apical and lateral borders of the disintegrated area until all of the pigmented cells are involved. The yolk cells retain their structure long after the pigmented cells have completely dis-




FIG. 5. View from animal pole showing stages in the disintegration of eggs in a four-cell stage after ten days' exposure to a temperature of 0° C. The eggs were in a two-cell stage when placed in the cold chamber.

integrated, except in the case of alcohol, which attacks the yolk rapidly from the start. Fig. 8 illustrates the general process.

(d) *Later Embryonic Stages.*—With the beginning of elongation, while two regions of high susceptibility are still present, viz., the apical and the dorsal lip region (now posterior), the posterior growing region, especially during the formation of the neural folds, appears relatively less susceptible than it did earlier. This apparent greater difference is probably due to local differences associated with the formation of the neural plate. In embryos beginning to elongate, disintegration begins first at the apical end in the medial dorsal region and spreads laterally and posteriorly, more rapidly in the posterior direction, to meet

an area of disintegration that has just begun in the region where the tail bud appears later. In neural fold stages disintegration begins at the anterior end of the floor of the neural groove usually at two points on either side of the median line, where the primordia of the optic vesicles have appeared. Often, and especially in earlier neural fold stages, the disintegration begins in the median line at the anterior end of the neural groove. From this




FIG. 6. Disintegration stages during late cleavage. The eggs are shown in side view with the animal pole uppermost and the gray crescent to the right.

point or points as the case may be, disintegration spreads back along the neural groove nearly to the posterior end, where it joins an area that has just begun to undergo dissolution. In the meantime, the medullary folds, ventral suckers, and anterior part of the head region have begun to disintegrate.

With the closure of the neural folds and the further elongation of the embryo, local differences in susceptibility begin to appear with the differentiation of certain organs. While in general, the apical-medial-dorsal region begins to disintegrate first, the dissolution of cells in this region may be followed shortly, or even preceded by, the disintegration of rapidly proliferating

cells in certain regions of the embryo, such, *e.g.*, as the tail bud, optic vesicles, nasal pits, ventral suckers, and other rapidly growing regions.

These data on disintegration are in agreement with the data on differential inhibition, acclimation, and recovery,¹ in showing that certain parts of the egg and embryo are more susceptible than other parts, and that these differences in susceptibility have a



FIG. 7. Camera lucida drawing of an egg showing disintegrated cells (white) following 24 hrs. exposure to $m/1,000$ KNC and 24 hrs. to $m/5,000$ KNC, from a two-cell stage. The animal pole is in the center of the figure. (Experiment KNC, B3.)

definite relation to the polar axis and to the plane of bilaterality in the early cleavage stages and to other physiological axes arising later in development. The significance of differential susceptibility is discussed later. (pp. 346-349).

2. *Experimental Modification of Development.*

For the sake of convenience in description, certain arbitrary terms have been used to designate different "types" of abnormalities. The differential inhibition of the cleavage ratio in early development $\left(\frac{\text{size of animal pole cells}}{\text{size of vegetative pole cells}} \right)$ is expressed as a fraction whose denominator is ten. For example, the expression "cleavage = $7/10$ " means that the size of the animal pole cells is to the size of the vegetative pole cells, as 7 is to 10. The expression "V-shaped blastopore" refers to a condition where the blastopore takes the shape of an inverted "V" or "U,"

and is intended to describe conditions where the dorsal lip is retarded to a relatively greater extent than the lateral lips (Fig 14, *A*). The term "wide-crescent blastopore" refers to a condition where the blastopore is in the form of a broad crescent (Fig. 17), and is intended to describe a condition where both dorsal and lateral lips are inhibited. The term "secondary invagination" describes a situation where a second infolding




FIG. 8. Stages in the disintegration of eggs in an early gastrula stage following several hours exposure to $m/1,000$ KNC, or in $m/1,000,000$ $HgCl_2$. The progress of disintegration follows this general plan in all agents in concentrations that kill the egg within several hours.

takes place apical to an original equatorial blastopore (Fig. 18). The "gastrular angle" is the angle between a plane through the center of the egg parallel to the floor of the segmentation cavity, and a line through the center of the egg and the blastopore (Fig. 9). Obviously, comparison of the gastrular angle of normal eggs with this angle in inhibited eggs is possible only at the *beginning* of gastrulation.

The development of the frog is so well known that a detailed description of the process is unnecessary. However, mention of several features of normal development will be of use for comparison with abnormal types.

Cleavage begins first and proceeds more rapidly in the pigmented hemisphere than in the vegetative hemisphere, establishing early in development an inequality in the size of the cells in the two hemispheres. In an early cleavage stage, the size rela-




FIG. 9. Diagram to illustrate the gastrular angle, *g.a.*

tion of the cells in the two regions is in the proportion of $4/10$ to $5/10$, a ratio that decreases in value up to about the time of gastrulation, when the ratio may be represented by $2.5/10$ to $3.5/10$. During mid-cleavage stages the cell size ratio is approximately $3.5/10$ to $4/10$. This does not take into account those




FIG. 10. *A*, control egg with a cleavage ratio of approximately $3.5/10$ - $4/10$. *B*, egg from the same batch as the one in Fig. 10, *A*, in which the cleavage ratio approximates $10/10$. Same treatment as egg illustrated in Fig. 7. (Experiment KNC B 3.)

pigmented cells on the grey crescent side of the egg, which are slightly smaller from the beginning of cleavage than the other pigmented cells (Morgan and Boring, 1903). Immediately preceding the appearance of the blastopore, those cells in the dorsal lip region are smaller than any of the other surface cells.

The gastrular angle at the time of the appearance of the blastopore approximates 40° .

In addition to the indications of bilaterality by the appearance of the grey crescent, bilaterality is made more apparent during the blastula stages by the movement of material from the pigmented region toward the equator—a process that takes place more rapidly in the sagittal plane than elsewhere, and results




FIG. 11. Enlarged segmentation cavity. Eggs in beginning two-cell stages treated as follows with KNC: $m/1,000$ 24 hrs., $m/5,000$ 24 hrs., $m/10,000$ 12 hrs., $m/20,000$ 12 hrs. (Experiment KNC H 5.)

FIG. 12. Small segmentation cavity as seen under conditions of severe inhibition. In this case eggs were exposed 48 hrs. from a two-cell stage to $m/10,000$ $HgCl_2$. (Experiment IV 40.)

in the thinning out of the walls and roof of the segmentation cavity, which now comes to lie nearer that side of the egg where the dorsal lip appears. The so-called germ ring, formed by the downward movement of animal pole material, and which lies




FIG. 13. Obliterated segmentation cavity. (Experiment IV 56.)

at first on the equator and later below it, extends somewhat farther down on the egg in the dorsal lip region. To express the situation in other words: growth in length (between the apical

pole and the germ ring) is greater and proceeds most rapidly in the sagittal plane. The significance of this process is discussed elsewhere (p. 349).

A. *Disturbances in Cleavage such that the Egg Never Gastrulates.*
 (a) *Cell Size Ratios.*—In frog eggs so strongly inhibited that they do not gastrulate, cleavage usually begins and proceeds more or less normally for several hours. The rate of cleavage in the animal hemisphere then becomes less and less rapid,




FIG. 14. V-shaped blastophore, A; Equatorial gastrulation, B; Secondary invagination, C. Produced in $m/500,000$ HgCl₂. About 40 per cent. of the eggs showed the V-shaped blastophore. (Experiment IV 70.)

relatively, until the cell size ratio approaches and may become equal to one ($10/10$, Fig. 10, B), whereas this ratio in a normal egg of a comparable stage of development is about $3.5/10$ to $4/10$ (Fig. 10, A).

The cleavage ratio was found to increase in value consistently in all cases¹ where inhibition was sufficient to prevent the com-

¹ This distortion of the cleavage ratio was obtained in: Experiments KNC A 1; H 1; exposure 12 hours in $m/1,000$ from 2-cell stage; cl. = $5/10$ to $7/10$. Experiment KNC—C.7; exposure 24 hours in $m/2,000$ from 2-4-cell stage; cl. = $7/10$. Experiment KNC—C.8; 48 hours exposure to $m/2,000$ from 2-cell stage; cl. = $8/10$. Experiment KNC—C.12; 48 hours exposure to $m/2,000$ from 2-cell stage; cl. = $7/10$ to $10/10$. Experiment KNC—C 1; 12 hours exposure to $m/5,000$ from 2-cell stage; cl. = $5/10$ to $6/10$. Experiment KNC—C.6; 24 hours exposure from 2-4-cell stage; cl. = $5/10$ to $7/10$. Experiment KNC—C.9; 5 days exposure to $m/10,000$ from 2 cell-stage; eggs beginning gastrulation (equatorial) with $6/10$ to $8/10$ cl. Experiment IV 21; 21 1/2 hours exposure to $m/8.5$ LiCl from an unsegmented stage; in about 15 per cent. to 20 per cent. cl. = $4/10$ – $7/10$. Experiment IV 31; unsegmented eggs exposed 4 hours in $m/4$ LiCl, then 7 1/2 hours in water showed an $8/10$ to $10/10$ cleavage ratio. Experiment IV 54; eggs in late cleavage exposed 40 hours to $m/100,000$ NgCl₂; cl. = $6/10$ to $10/10$. Experiment IV 62; eggs in 4-8-cell stage exposed 12 hours in 0.0075 per cent. formaldehyde; cl. = $6/10$ to $10/10$. Citations of experiments in italics are from Dr. Child's preserved material. "cl." = cleavage ratio.

pletion of gastrulation, although the proportion of eggs showing this type of differential inhibition varied somewhat in experiments involving different chemicals. This increase in value of the cleavage ratio appears less frequently in LiCl, MgCl₂, and in alcohol, and most frequently in KNC and formaldehyde. In $m/1,000$, $m/2,000$ KNC, fully 90 per cent. of the eggs are of this type after 24 to 36 hours' exposure from the beginning of segmentation. Differential inhibition under the conditions stated, is diagrammatic, especially in KNC, and represents simply a greater susceptibility to the toxic agent, of the cells most affected, viz., the animal pole cells. The significance of this differential susceptibility is a matter of discussion elsewhere (pp. 346-349).

(b) *Bilaterality*.—Under the conditions of these experiments, bilaterality is much obscured and may not become evident at all (Fig. 20). But, that bilaterality is at least potentially present is indicated in those cases where gastrulation does begin (always equatorially) the wrinkling and folding, indicating the initiation of gastrulation usually begins at a definite point on the equatorial region and spreads equatorially around the egg. A few cases have been observed in $m/5,000$ KNC where the equatorial folding occurred almost simultaneously around the entire equator of the egg.

(c) *The Segmentation Cavity*.—Under severe inhibiting conditions, the segmentation cavity shows several characteristic consequences of differential inhibition, depending chiefly upon the severity of the conditions and the stage at which the eggs are exposed to them and upon the length of the exposure.

Some of the data are these: In eggs exposed to $m/1,000$ or $m/5,000$ KNC (Experiments KNC A-H) from the beginning of the first segmentation, development usually ceases in late cleavage stages with occasional abortive attempts at equatorial gastrulation. The yolk cells, which are relatively less inhibited, may continue division for a time after it has nearly or quite stopped in the animal hemisphere. In such cases, one finds that the more peripheral yolk cells, forming the floor of the segmentation cavity, proliferate apicalward, forming a thin layer of yolk cells that may partly or completely line the walls of the segmentation

cavity—which becomes very large. At the time of death, the egg resembles in some respects an amphioxus blastula (Fig. 11). Superficial indications of bilaterality are obscure.

Where inhibition is more severe, *e.g.*, in $m/10,000$ HgCl_2 (Experiment IV 40), eggs exposed to the solution for 48 hours from a two-cell stage, stop development in late cleavage stages. Gastrulation never occurs. In all of the eggs, the downward migration of materials from the apical pole is completely inhibited and the walls of the very small segmentation cavity lying near the center of the egg, are $2/5$ to $4/9$ the diameter of the egg in thickness (Fig. 12).

The partial or complete obliteration of the segmentation cavity may occur in an entirely different way under less severe inhibiting conditions. Eggs in late segmentation stages, exposed 15 minutes to $m/10,000$ HgCl_2 (Experiment IV 56) then returned to water, show after three days, the segmentation cavity completely filled with small yolk cells that have proliferated from the floor of the cavity. A faint line indicates the walls of the now obliterated cavity. The walls of what was the segmentation cavity are much thinner than when the eggs had been exposed 48 hours to $m/10,000$ HgCl_2 from the beginning of cleavage.

Likewise in $m/10$ LiCl , where development proceeds abnormally to the time of hatching, the proliferation of yolk cells from the margin of the floor of the segmentation cavity may be so extensive as to fill it completely.

All of these different modifications appear to be simply different expressions of the differential inhibition which the eggs have suffered. The animal pole cells being most susceptible, are most affected by the adverse conditions, the yolk cells least.

Those small cells around the margins of the floor of the segmentation cavity, which under certain conditions of inhibition, continue division and finally fill the blastocoel completely, appear to be the ones destined to give rise to mesoderm (Morgan, 1906, p. 129; Kellicott, 1913, p. 107; King, 1902, Fig. 4). It is significant here that in the sea urchin Child (1916, p. 91) found the mesenchyme cells which arise from the basal pole of the egg, to be less susceptible than other parts of the egg, and that under conditions of differential inhibition, these cells tended

to "run wild" and result in an excessive over-development of skeletal structures.

(d) *Meroblastic Cleavage*.—In exceptional cases, and under the influence of conditions so severe that development stops completely in early cleavage, cell division may be partly or completely restricted to the animal pole. This type of cleavage has been observed following exposure: to urea, 2.34 per cent. (Jenkinson, 1906, Figs. 38, 41); to NH_4I (Jenkinson, 1906, Fig. 32); to Na_2SO_4 (Jenkinson, 1906, Fig. 40); to temperatures above 26°C . (Hertwig, 1895), et al. In my own experiments it has been observed among eggs exposed six to ten hours to 2 per cent., 3 per cent. alcohol, a result that is not surprising when one considers the solvent action of alcohol upon lipid substances generally. Four factors seem primarily concerned in the production of these restricted cleavages, viz., time, concentration or intensity of action of the agent used, physical effect of the particular agent to which the eggs are exposed, and the stage at which the eggs are exposed to the agent. So far as my observations go, this type of cleavage, rare at best, is realized only when the eggs are exposed to the inhibiting conditions at the time of, or immediately preceding the appearance of the first cleavage plane. If the eggs are exposed to concentrations necessary to the production of this type of cleavage, several hours before cleavage would normally begin, they never segment. The physical effect of alcohol has been mentioned. The time element is more or less obvious. Cleavage begins first, *normally*, in the pigmented hemisphere, and under the conditions necessary for the production of meroblastic cleavage the first cleavage plane makes its appearance, in all probability, before the effect of the inhibiting conditions (in the case of chemical substances particularly) penetrate the gelatinous membranes of the egg.

B. *Disturbances in Gastrulation*.—The various modifications of the process of gastrulation, following inhibition, fall into several more or less distinct groups, the particular types obtained depending largely upon the severity of the inhibiting conditions and upon the stage at which the eggs are introduced into the experiment; *i.e.*, depending upon the treatment and the physiological condition of the eggs. Individual variation is an impor-

tant limiting factor in the experimental control of modifications. The range of variation in this respect seems to be widest during the early cleavage stages. In the great majority of experiments, while one usually finds a complete series of stages from nearly normal to the most extreme deviations from the normal, the modifications produced tend to conform to certain types, varying with the physiological condition of the eggs and the treatment they receive. It is therefore necessary to speak in terms of averages in discussing a given type of abnormality.

In general, as the concentration of a given chemical to which the eggs are exposed, is increased, or with longer exposure to the same concentrations, the modifications produced become more extreme. They are discussed in the order in which they appear.

(a) *Retardation of the Dorsal Lip Region.*—These are stages




FIG. 15. Inhibition of the dorsal lip region after 38 hours exposure to $m/10.62$ LiCl. Eggs placed in the solution were in 16-32-cell stages—24 hours after deposition. (Experiment IV 76.) *d.l.*, dorsal lip region; *y.p.*, yolk plug.

in which only the dorsal lip region seems much inhibited, although occasionally the cleavage ratio may increase in value somewhat, and the gastrular angle (see p. 329) in a small per cent. of the cases may be slightly more acute than in the control eggs. Where only the dorsal lip region is inhibited, the blastopore takes the form of an inverted V or U and when complete the yolk plug is oval, and may be several times longer than broad in the sagittal plane.

Experiments IV 70; G, H, I. Eggs deposited March 28, 3:00 P.M. Removed to the laboratory 4:00 P.M. Introduced into $m/500,000$ HgCl_2 , 5:00 P.M.—unsegmented at this time. After 24 hours the eggs were washed and returned to water. Twenty-four hours later (March 29, 5:00 P.M.) the stages shown in Fig. 14 were recorded. About 40 per cent. of the eggs showed the V-shaped blastopore (Fig. 14, A). Eggs in the three experiments were alike and among them everything was present from a few with closed blastopores to several forms showing a secondary invagination midway between the original (equatorial) blastopore and the apical pole. The equatorial gastrulae resemble those produced in $m/10$ LiCl.

Experiment IV 91. Unsegmented eggs, placed in $m/2,000,000$ HgCl_2 . Forty hours later, the eggs were mostly in inhibited gastrula stages (V-shaped blastopores).

Eggs introduced into $m/10.62$ LiCl during 16–32 cell stage, exhibit the inverted V- or U-shaped blastopore after 24 hours' exposure (Experiment IV 76). Eggs introduced into $m/10$ LiCl immediately before gastrulation (Experiment IV 59) are in gastrula and large yolk plug stages after eighteen hours in the solution. The majority of gastrula stages show the long inverted V-shaped blastopore and in those eggs showing yolk plugs, there is a tendency for the yolk plugs to be much larger than normal and somewhat oval in shape (Fig. 16).

After 48 hours in the solution, ninety per cent. of the eggs were in early neural fold stages, with much elongated yolk plugs.

In experiment IV 62 unsegmented eggs, five hours after deposition, were introduced into 0.0075 per cent. formaldehyde, and varied from equatorial gastrulae to early neural fold stages after sixty hours' exposure. A few showed secondary invaginations.

In experiment R 138 a, eggs in early gastrula stages were placed in $m/1,000$ KNC. After a 12-hour exposure, 90 per cent. of the eggs showed a long inverted V-shaped blastopore or much elongated yolk plug. The eggs did not develop farther in this solution.

(b) and (c) *Retardation of Both Dorsal and Lateral Lips of the Blastopore, and Equatorial Gastrulation.* Eggs showing retardation of both dorsal and lateral lips are described as "flat-crescent"

gastrulæ, for the reason that under the slightly more severe inhibiting conditions where both dorsal and lateral lips are inhibited, the blastopore forms, when about half completed, a figure like a much flattened crescent. When such a blastopore




FIG. 16. Inhibition of dorsal lip region. Eggs placed in $m/10$ LiCl in late cleavage stages immediately before gastrulation. A and B after 18 hrs. in the solution; C, D and E after 54 hrs. in the solution. *d.l.*, dorsal lip region; *y.p.*, yolk plug. (Experiment IV 76.)

is complete the yolk plug is always much larger than in the control. All stages exist between this flat crescent type and those cases where the blastopore extends around the equator of the egg (equatorial gastrulation). In these forms the cleavage ratio

varies from nearly normal (in the less inhibited ones) to 10/10 (more inhibited ones). In the more extreme inhibitions some cases of secondary invaginations are seen and a decrease in the gastrular angle is characteristic. Almost any type may be produced at will simply by varying the concentration of the inhibiting agent, or the length of the exposure, or the stage at which the egg is introduced into the experiment, or by varying several of the factors at once. The chief limiting factor in the control of these and other modifications is the variation in susceptibility exhibited by the eggs of different females. Several preliminary attempts may be necessary before the desired result is obtained. Some of the experiments are cited below, in which these modifications were prominent, and which serve to illustrate a few of the many different ways in which these abnormalities may be produced.

Experiment IV 21.¹ Unsegmented eggs, introduced into




FIG. 17. Inhibition of both dorsal and lateral lip regions of the blastopore, resulting in a "flat-crescent" blastopore. 28 1/2 hours' exposure to $m/10.62$ LiCl from unsegmented stage. (Exp. IV 22.)

$m/7$ LiCl 1 1/2 hours after fertilization. After 21 hours in LiCl and 7 hours in water a few of the eggs showed a slight equatorial wrinkling but no distinct blastopore was present. After 24 hours in LiCl, and 24 hours in water most of the eggs were in late blastula stages in which the segmentation cavity was completely lined with one or more layers of small round slightly pigmented cells.

Experiment IV 21. After 20 hours in $m/8.5$ LiCl and 8 hours

¹ Eggs used in experiments IV 21, 22, 26, 27, 29, 31, 33, 34, all came from the same female, and were introduced into the different experiments 1 1/2 hours after deposition.

in water, about 80 per cent. of the eggs were beginning equatorial gastrulation. After 20 hours in this solution and 22 hours in water, about 70 per cent. of the eggs showed various stages of equatorial gastrulation; the rest of the eggs varied from nearly normal yolk plug stages to those showing flat-crescent blastopores (Fig. 17). After 76 hours in water following 20 hours' exposure




FIG. 18. Secondary invagination following 48 hours' exposure to $m/10$ LiCl, from an unsegmented stage. A, surface view; B, sagittal section of same egg. *inv. 2*, secondary invagination; *d.l.*, dorsal lip; *bp.*, blastopore; *s.c.*, segmentation cavity. (Exp. LiCl 1 d.)

to the solution about 5 per cent. of the eggs showed a secondary invagination apical to the equatorial blastopore. This secondary invagination appeared more prominently (in about 40 per cent. to 50 per cent.) after 28 hours in $m/8.5$ LiCl and 48 hours in water (Fig. 18).

Experiment IV 22. 48 hours' exposure to $m/10.62$ LiCl gave 90 per cent. equatorial gastrulae (Fig. 19), which when returned to water proceeded to develop farther and after 28 hours were in elongated neural fold stages with large protruding yolk plugs. After 72 hours in water following 48 hours' exposure to the solution one finds all stages from a few equatorial gastrulae to microcephalic and anencephalic forms with partially or completely "fused" suckers, nasal pits, and optic vesicles. After 76 hours' exposure to the solution and 20 hours in water the abnormalities were more extreme. In about 5 per cent. of these the pigmented cells grew out from one to ten mm. to form a cone of cells radially symmetrical about the original polar axis of the egg, there being

not the slightest external evidence of bilaterality or dorso-ventrality (Fig. 20).

Unsegmented eggs exposed 48 hours to $m/1,000,000$ $HgCl_2$ exhibited all stages between equatorial gastrulation and fairly




FIG. 19.

FIG. 20.

FIG. 19. Equatorial gastrulation following 76 hours' exposure to $m/10.62$ $LiCl$ from an unsegmented stage. (Experiment IV 22.3.)

FIG. 20. Embryo radially symmetrical about the polar axis. Produced by exposing unsegmented eggs 76 hrs. to $m/10.62$ $LiCl$; 20 hrs. in water. (Experiment IV 22 b.)

normal yolk plug stages. Eggs with a broad crescentic blastopore were most prominent. The eggs died without developing farther.

Experiment IV 34. Eggs in 8-cell stage (10 hours and 20 minutes after deposition) were placed in $m/5,000$ $KMnO_4$. After 48 hours' exposure—solution not changed—the eggs showed a complete series of stages from perfect equatorial gastrulae to normal yolk plug stages. At this time all of the permanganate had been reduced to the brown form and the solution had evidently lost much of its toxicity. Sixty hours after being placed in the solution about 50 per cent. of the embryos were elongating but were markedly macrocephalic, showing differential recovery.

(d) *Secondary Invaginations*.—The eggs described by this term show a secondary invagination appearing in a meridian bisecting the original blastopore which is usually equatorial. The invagination may extend partly or completely around the egg and

appears only under conditions of severe inhibition. In addition to this secondary inturning, some overgrowth may take place; in some cases the over growth is in the direction of the vegetal pole as in normal gastrulation, while in other cases the overgrowth is toward the apical pole. In eggs where this upward




FIG. 21. Some types of modification seen after the following treatment in KNC: A, B (same egg). Eggs beginning two-cell at start of experiment; 24 hrs. in $m/1,000$, 24 hrs. in $m/20,000$, 12 hrs. in $m/50,000$, 12 hrs. in $m/100,000$. C, 12 hrs. $m/1,000$, 12 hrs. $m/5,000$, 24 hrs. $m/10,000$. (Experiment KNC A 3.) Note the position of the blastopore which is nearer the equator than in control eggs.

overgrowth is taking place over the secondary invagination, the yolk cells below the original blastopore are usually overgrowing the pigmented cells of the animal hemisphere, thus reversing the usual process.

This modification is similar to that described by Morgan (1903), which he says is produced by a process in which "the cells of the upper hemisphere . . . turn into the egg." It makes little difference whether one refers to the process thus or whether one calls it simply an invagination. I have used the latter term because the process appears to me to be similar in many respects to that of invagination and overgrowth seen in gastrulation.

Some of the experiments in which this secondary invagination appeared have been cited. They are illustrated in Figs. 14, C, and 18. Several other experiments follow.

Experiment LiCl 1. Unsegmented eggs placed in $m/10$. 6 hours: not appreciably different from control. 24 hours: beginning equatorial gastrulation. Pigment has decreased somewhat in the cells of the dorsal lip region. The probable significance of this disappearance of pigment has been mentioned (p. 321). 30 hours: large yolk plug—equatorial gastrulae. 48 hours: equatorial gastrulae with secondary invaginations.




FIG. 22. A, differential inhibition—fused suckers, nasal pits, dorsal concavity. Three hours' exposure (from late cleavage) to $m/5$ LiCl; four days in water. (Experiment IV 58 b.) *v.s.*, ventral sucker; *n.p.*, nasal pit. B, differential recovery. Note the marked dorsal convexity. (Experiment IV 59 e.) Two days' exposure (from late cleavage) to $m/10$ LiCl; five days in water.

In experiment IV 70 (see p. 337) this modification occurred in about 10 per cent. of the eggs following 24 hours' exposure to $m/500,000$ $HgCl_2$ and 24 hours in water. A few cases were noted following 48 hours' exposure, from unsegmented, to 0.0075 per cent. formaldehyde (Experiment IV 83). In experiment R 135, eggs just beginning to gastrulate were placed in 0.001 per cent. formaldehyde and after 24 hours' exposure showed equatorial gastrulae with secondary invaginations similar to the one shown in Fig. 18. Eggs just beginning to segment when subjected to the following graded treatment showed various degrees of this modification in about 50 per cent. of the eggs: KNC $m/1,000$ 24 hours, $m/20,000$ 24 hours, $m/50,000$ 12 hours, $m/100,000$ 12 hours. (Experiment KNC H 5).

(e) *The Gastrular Angle*.—In the normal egg the blastopore normally makes its appearance in a meridian that bisects the grey crescent and about 40° below a plane through the center of

the egg and parallel to the floor of the segmentation cavity. It is characteristic of strongly inhibited eggs that the gastrular angle approaches and often equals 0. Not infrequently the angle becomes negative, *i.e.*, a circular blastopore is formed *above* the equator. Some of the experiments where this shifting of the point of appearance of the blastopore occurred have been cited (pp. 337, 339, 340). Several variations of this modification are illustrated in Fig. 21.

Under conditions where marked acceleration of development occurs, *e.g.*, in $m/100,000$ KCN; $n/5,000$ to $n/20,000$ HCl⁶ (2 c.c.




FIG. 23. A, normal. B, microcephalic, eyes closer together than normal, nasal pits fused, ventral suckers absent. Three hours' exposure (from early gastrula) to $m/7$ LiCl; 13 days in water. (Experiment R 122.1.)

to 5 c.c. of $n/10$ HCl in well water) made up in well water, there is a tendency for the gastrular angle to approximate 45° or even 50° . It is much more difficult to alter the space relation of the blastopore and apical region by acceleration than by inhibition. It may be noted that the effects of acceleration become more apparent about the time of hatching. In these cases the embryos are markedly macrocephalic and dorsally convex, indicating more rapid growth of the anterior and dorsal parts.

C. *Differential Inhibition in Later Development.*—The whole problem of differential susceptibility involves of course, not only differential inhibition, with which this paper is primarily concerned, but it also includes differential acclimation and recovery,

⁶ I am aware that acid is generally regarded as an inhibitor of protoplasmic activity, but in these experiments the acceleration of development in certain concentrations of HCl is perfectly definite and distinct. Similar results were obtained independently and repeatedly by both Dr. Child and myself. Since the well water in which the solutions were made up contains various salts and other substances in minute amounts it remains to be decided whether the acceleration of development in these cases is a direct or an indirect effect of the acid.

the data on which are fully as striking and significant as those on differential inhibition. But their consideration is reserved for future consideration along with a detailed discussion of differential susceptibility in the later stages of development in



FIG. 24. Cyclopic frog embryo. (Experiment R 122.1.)

the frog's egg. However, for the sake of completeness a few of the more striking abnormalities of later development may be mentioned. These are: microcephalic forms with such bilateral structures as optic vesicles, ventral suckers, and nasal pits in all degrees of approximation from nearly normal to complete "fusion."¹ These forms are seen somewhat more frequently in




FIG. 25. Spina bifida, microcephalic, ventral suckers nearly fused. Twelve hours' exposure (from early gastrula) to $n/300$ NaOH; four days in water. (Experiment R 125.4.)

$m/10$ LiCl than in other agents but they have been obtained in varying percentages in all of the agents used to modify development. Several of the forms are illustrated in Figs. 22-25. The history and treatment of the eggs is given in the legends to the figures.

In addition to the marked apical inhibitions just mentioned the later stages show also considerable median dorsal inhibition

¹ The term "fusion" is used in a purely descriptive sense for the eye of cyclopic forms, single median nasal pits, or single median ventral suckers. Whether the primordia of the organs mentioned arise as paired structures and later fuse to form a single median structure, or whether a single primordium arises that later divides into two parts in normal individuals, but which do not separate under strongly inhibiting conditions, is a question for future consideration.

(Fig. 22, *A*) indicated by the marked dorsal convexity with the tail extending upward, in some cases at right angles to the body. The medullary folds may fail to close in the brain region, in the brain region and at the posterior end, or they may not close at all throughout their entire length. In some cases anencephalic forms appear. Spina bifida of all degrees is of course common under conditions that inhibit development, and result primarily from inhibition of the dorsal lip region.

V. DISCUSSION.

It has been shown that gradients in susceptibility to several external agents do exist in the egg and embryo of frog. Concerning the *fact* there can be no question. It was pointed out that the experimentally produced modifications described in this paper are readily and logically accounted for on the basis of differential susceptibility. Now, since no evidence was found of: (*a*) "specificity" in the sense in which the term has been used to explain terata produced by various chemical agents; (*b*) of "blastolysis" or (*c*) of any "disorganization" effect of the chemicals used to modify development as effective factors in the production of terata, the interpretation of the data is now largely a matter of pointing out the basis and significance of differential susceptibility. This is the chief task of the discussion.

1. *The Nature of Physiological Axes.*—These differences in susceptibility to external agents exhibited by different parts of the living egg and embryo (see pp. 322-344) must represent differences of some sort in protoplasmic conditions. Living protoplasm is a complex system of correlated, interdependent reactions and conditions, and it is unquestionably true that different agents affect it differently or attack it at different points. But, whatever the particular factor or factors chiefly concerned in determining these differences in susceptibility to particular agents may be, whether it be rate of fundamental oxidation-reduction processes, state of colloidal aggregation, permeability of limiting membranes, or something else, the general uniformity and lack of specificity in the susceptibility relations to different agents and conditions, not only as regards the egg and embryo of the frog but as regards animals and plants generally, force us to

certain conclusions. These are briefly: (1) that differences in susceptibility must depend primarily upon quantitative differences in protoplasmic condition, rather than upon the specific or qualitative differences, and (2), that these quantitative differences are closely associated with differences in the rate at which fundamental physico-chemical processes are taking place in different regions of an individual or in different individuals.

These facts, viz., differential susceptibility relations and their marked uniformity and lack of specificity, not only in the frog but also in other organisms; gradients in protoplasm-yolk and pigment distribution both of which parallel the polar axis; and the fact that development begins and proceeds in a definite, orderly, and sequential manner with respect to certain axes (or to the polar axis) in such a way that structural and functional order are at all times referable to one or more axes or planes of symmetry (anterior-posterior, dorso-ventral, medio-lateral), constitute evidence for the existence of a fundamental underlying order of some sort that exists prior to, and which conditions the orderly, sequential, and spatial relations of parts that become apparent later in development. Furthermore it is this order, of which differential susceptibility is one expression, which affords the basis for the great uniformity and lack of specificity in the susceptibility relations, not only in the frog but also in other organisms.

The nature and origin of this underlying order is the problem of physiological axiation—a problem that has been the subject of no little speculation and experimentation. It has been thought by some that polarity and symmetry as manifested in the organism are fundamental properties of protoplasm, and, like the polarity and symmetry of crystalline substances or of the magnet, are intra-molecular in character. It may be said, however, that this conception and others of a similar nature, do not stand the test of experiment even where capable of such an analysis.

On the basis of extensive experimental work Child has arrived at a dynamic conception of physiological axes in the organism with the aid of which he accounts in a satisfactory manner not only for many of the hitherto unrelated facts of normal and abnormal development, but also he has been able experimentally

to modify and control development on a predictable basis.¹ Child looks upon a physiological axis in its simplest terms as a quantitative gradation in physiological condition. He has brought forward evidence from a number of sources which point to the conclusion that this gradient is a gradation in rate of metabolism, or at least of certain fundamental reactions such as the oxidation-reduction processes, which is associated with a gradation in the physical condition and constitution of the protoplasm.

If a gradient in physiological condition is a fundamental ordering factor in the process of physiological axiation, it should be possible to obtain as many expressions of this gradient as there are aspects into which fundamental metabolic activity and the physical constitution of the protoplasm may be analyzed. Hence, we might expect to find, *e.g.*, gradients in irritability; in growth and differentiation and development; in the visible morphological condition of the protoplasm; in permeability; in the active mass of enzymes; in electrical potential; in oxidation as measured by oxygen consumption and in carbon dioxide production; and conversely, gradients in susceptibility to lack of oxygen. Well known biological facts afford demonstrative evidence for the existence of some of these gradients. Further evidence for the existence of other of these gradients has been published by Child, Hyman, et al.¹ Other evidence is yet unpublished.

"All the various lines of evidence considered agree in showing that axial gradients in the dynamic processes are characteristic features of organisms and that a definite relation exists in each individual between the direction of the gradient in any axis and the physiological and structural order which arises along that axis. In the major axis the region of highest rate in the metabolic gradient becomes the apical or anterior region of the individual, and in the minor axes also the regions of highest rate in the gradients represent particular features of the order in each case. Along any axis particular parts apparently represent particular levels in the gradient. The variety, extent, and agree-

¹ See footnote 1, page 315.

ment of the evidence is all the more interesting in view of the fact that such gradients have not heretofore been recognized as characteristic features of organic constitution." (Child, 1915*a*, p. 87.)

2. *Origin of the Dorsal Lip Region.*—The data on differential killing and differential inhibition in the frog show, not only that the apical pole region is more susceptible than other parts of the egg during early development, but also that as development proceeds a secondary region of high susceptibility appears in the sagittal plane about 100° below the apical pole. This is the dorsal lip region. Morgan and Boring (1903) showed that in the eight cell stage, that cell of the upper quartet lying nearest the grey crescent is slightly smaller than the other three cells of the quartet. This inequality persists up to a stage some time after the beginning of gastrulation, cell division being somewhat more rapid here than elsewhere on the egg. It is well known that during early cleavage some of the apical cells migrate or are pushed downward to form a so-called germ ring that lies at first on the equator of the egg and later below it. Correlated with the greater activity of the apical pole cells in the sagittal plane, this migration or pushing is most rapid in this plane, *i.e.*, in that pigmented quadrant the median meridian of which bisects the grey crescent. This displacement of material from the apical pole region is probably not to be conceived as an active migration but as a consequence of the more active cell division and growth in the embryonic meridian. After the materials in this meridian have been displaced a certain distance from the apical pole they seem to get beyond the range of correlation of that region; certainly they become more active and establish the posterior growing region (dorsal lip). At the time of gastrulation some of the material included in the dorsal lip cells is certainly not less than 100° from where it was in the four-cell stage. What this migration of the materials of the apical region toward the equator of the egg amounts to, of course, is a growth in length which is greatest and proceeds most rapidly in the sagittal plane.

This dorsal lip region, which arises secondarily as a rapidly growing posterior region, appears normally at a fixed distance (about 130°) from the apical end of the polar axis, and it is

shown that this distance decreases with increasing inhibition (decrease in metabolic rate) of the apical region, and increases slightly with increase in activity (acceleration) of the apical region. The facts suggest at once a definite physiological correlation between the apical region and the region of the dorsal lip which arises secondarily as a posterior region of active growth and differentiation, and which does not appear to differ fundamentally from the posterior growing region which gives rise to the trunk in annelid larvæ (Child, 1917a¹).

From these facts, viz., the secondary origin of the dorsal lip region, its definite spatial relation to the apical region, which may be controlled experimentally, and its appearance at the distal end of a rapidly growing region, the origin of the dorsal lip region offers an interesting parallel to those cases in certain lower animals and plants where new zooids or new individuals arising by asexual reproduction have been shown to appear as physiological isolations.¹

In the simple cases of physiological isolation in lower animals

¹ The rôle of physiological isolation, dominance and subordination in development has been studied by Child and fully stated by him (1915a, pp. 88-169) and need not be considered here except to mention some of the pertinent facts. In Child's own words:

"The experimental evidence demonstrate, first the essential independence of the apical region in both plants and animals, and secondly, determination and control by this apical region of the developmental processes at other levels of the major axis of the individual." (1915a, p. 125.)

The facts indicate for those forms investigated, that the control of subordinate parts by the dominant region is transmissive rather than transportative in character, and that the range of dominance, *i.e.*, the distance over which the control of the dominant region is effective and beyond which physiological isolation occurs, varies with the relative metabolic rates of the dominant and subordinate parts, and condition of the conducting path; increasing with increase in metabolic rate in the dominant regions and with the specialization of the conducting path, and decreasing with decrease in the metabolic rate of the dominant region, with increase in rate of the subordinate region, and with interference with the conducting path. These four ways in which physiological isolation may occur are summed up by Child, thus:

"It is possible to control and alter experimentally the spatial relations of parts in the individual by altering the length of the metabolic gradient and so the range of dominance. Parts of the individual may come to lie beyond the range of dominance in consequence of increase in size of the whole, of decrease in range and degree of dominance by decrease in the metabolic rate in the dominant region, of decrease in conductivity of the paths of correlation, and of the direct local action of external factors which increase the independence of subordinate parts." (1915a, p. 169.)

such, *e.g.*, as the flatworms, it appears that as the body increases in length the posterior region comes to lie beyond the range of dominance (control) of the apical end. As a consequence, the posterior end undergoes much the same changes that it would if physically isolated. It begins to dedifferentiate and its susceptibility increases, *i.e.*, it comes to resemble physiologically younger tissue. If these processes go far enough the posterior region gives rise to a new zoöid which in time may become a new individual.¹

In the annelids such a region arises early in development but instead of producing zoöids which develop into independent individuals it proceeds to give rise to segments that are reintegrated and brought under control, probably through the specialization of the nervous system. From this point of view the development of the annelid trunk appears to be a succession of incomplete reproduction processes, but which undergo reintegration into a whole consisting of segments.

In the light of all the facts it is difficult to avoid the conclusion that the origin of the dorsal lip region is fundamentally similar to these processes of physiological isolation, since the dorsal lip region arises secondarily at the posterior end of the embryonic area as a region of high susceptibility, increased activity, and rapid growth.

The secondary invaginations afford an interesting bit of evidence in that, with certain degrees of inhibition of the apical region, the range of dominance is decreased and after a first isolation has occurred—a second one may arise nearer the apical end.

3. *The Question of "Specificity" in Teratogeny.*—The word "specificity" as used in the literature on vertebrate teratogeny is difficult of exact definition. In general it has been used to designate some unique reaction of an organism or part, to a particular experimental factor, especially a particular chemical substance, such as to produce an anomaly characteristic of that individual external factor.

Various attempts have been made to identify experimentally produced abnormalities with either a specific chemical effect or with a more general physical effect of some one of numerous

¹ Child, 1911, *Jour. Exp. Zoöl.*, Vol. 11.

agents that have been used to modify development. The literature of this subject has been reviewed incidentally so many times (Korschelt and Heider, 1902; Hertwig, 1906; Jenkinson, 1909) that only certain essential facts and conclusions need be mentioned here. Gurwitsch (1896) attributed the various abnormalities of the frog produced by him in sodium bromide, lithium chloride, and weak solutions of strychnine, caffeine, and nicotine, to an "einigermaassen spezifische" action of individual chemicals upon certain regions of the egg. Stockard (1909) at one time believed that cyclopia in *Fundulus* was due to a specific cyclopic producing property of $MgCl_2$ and went so far as to suggest that cyclopia in man may be due to an unusually high amount of Mg in the amniotic fluid or in the blood of the pregnant mother. When, however, it had been shown that a large number of different substances produce the same anomaly, Stockard (1914) abandoned his earlier view that there are "specific responses to the given chemical substances employed" and suggested that since "a certain definite response on the part of the developing organism may be consistently obtained after carefully adjusted treatments with a large number of different substances . . . in certain cases they may serve simply to lower the developmental metabolism and thus prevent or arrest the formation of particular structures." No reason is offered to explain *how* lowering the developmental metabolism may cause the prevention or arrest of the formation of particular structures or *why*, under properly controlled conditions, *particular* regions of the embryo are affected more than other parts and in a definite and sequential relation to one another, though he notes that the nervous system and special sense organs are most affected by the use of chemical agents that inhibit development.

To digress for a moment, in the light of present knowledge we believe it possible to give a rational interpretation of these facts. The reason that a lowering of the developmental metabolism under conditions where acclimation does not occur, can prevent or arrest the formation of such structures as nervous system and special sense organs, is the fact that there is a gradient in the rate at which the processes described by the collective term "metabolism" are taking place along physio-

logical axes. Anterior, medial, dorsal regions, because of their higher metabolic rate are more susceptible to the inhibiting conditions and hence are more affected.

Bataillon (1901) studied the abnormal development of the frog in isotonic solutions of cane sugar, NaCl, and a number of other salts and reached the conclusion that osmotic pressure and the consequent withdrawal of water from the developing embryo, is the effective cause of abnormalities. Jenkinson (1906) showed that solutions of a large number of substances, isotonic with 0.625 per cent. NaCl, do not produce the same, but markedly different effects. It may be pointed out however that these effects differ not in kind, but in degree, and are on the whole readily interpreted in terms of differential susceptibility. And it may be remarked here that the large number of abnormalities in the frog described during the past quarter of a century which have been produced by experimental means may all be rationally interpreted in these terms and readily brought into relation with the axial gradients. Jenkinson's conclusion was that the abnormalities produced—covering most of the modifications known in amphibian teratology—were not to be attributed to the osmotic pressure of the solutions but were due to their chemical or physical properties. Morgan who has done much work on abnormal development in the frog, stated, (1906) in speaking of the osmotic and chemical effect of salt solutions on the frog egg, that: "It is probable that the effect is a double one; in part chemical, in part osmotic."

There can be little doubt that certain substances especially in high concentrations, do have a physical as well as a chemical effect upon protoplasm. Furthermore, in a highly specialized egg like that of the frog where a considerable volume of relatively inert matter (yolk) is localized in one hemisphere, high osmotic pressures such as exist in solutions of certain electrolytes and other substances in concentrations high enough to cause markedly abnormal development, and substances that have a solvent action on the yolk (alcohol, *e.g.*), may in extreme cases set up secondary disturbances of a mechanical sort such as partly or wholly to obscure the general susceptibility relations. But, it is especially to be noted that those agents which, in extreme concentrations

or intensities of action, do tend to obscure the differential susceptibility relations, may be used in other concentrations or intensities, to produce differential inhibitions that, in kind, are not different from the differential inhibitions produced in a variety of other agents.

Although it was stated (p. 346) that in the reaction of the frog egg to the different substances used to modify development, no evidence was found of any "specific" action, *in the sense in which the term has been used*, I do not wish to be understood as implying that all chemical substances act upon protoplasm in exactly the same way. It is to be expected, and bio-chemical and zoölogical literature contain abundant evidence to show, that different agents do affect protoplasm in different ways. KNC, *e.g.*, which is a powerful reducing agent is known to interfere with oxidations and enzyme activity in the organism. Hyman (1919) found that in planaria oxygen consumption may be reversibly cut down as much as 80 per cent. to 90 per cent., in the presence of minute amounts of KNC. In this paper she gives a comprehensive review of the literature on the action of cyanides on protoplasm so that further discussion of this matter is unnecessary here. So far as susceptibility relations are concerned formaldehyde appears to affect protoplasm in much the same way as KNC, and KMnO_4 which is a vigorous oxidizing agent is equally efficient for demonstrating differences in susceptibility in organisms.¹ It is a significant fact that these oxidizing and reducing agents are among the most effective agents that have been used in the study of physiological axiation.

As regards anesthetic action in the stricter sense, whether these agents disturb oxidations, affect the lipins of the cell, produce changes in the surface membranes, etc., there appears to be no consensus of opinion.

Various salts affect protoplasm, especially the colloid substances, in widely different ways. They may bring about changes in the state of aggregation of colloids, and the loss or addition of water; induce changes in ionization which may result in the precipitation or solution of certain substances in the protoplasm; or upset the electrical equilibrium, and so on.

Acids and alkalies also affect the speed and type of many

reactions in protoplasm in a variety of ways depending upon the ionization constant of the acid or alkali, concentration, reactivity of the substances formed by their interaction with protoplasm, etc.

It is obviously impossible here to go into a detailed discussion of the particular types of reaction set up between external agents and living protoplasm. For a resume of this important subject the reader is referred to Mathew's "Physiological Chemistry," especially chapter V., where references to the literature may be found.

In spite of the various factors concerned in the action of external agents upon living protoplasm, the highly significant fact remains that the susceptibility relations to external agents whose action is severe enough to kill within a few hours or to be distinctly toxic show not only a high degree of uniformity for at least a large number of different agents, but also a very definite correlation with metabolic conditions.

As regards permeability of limiting membranes there can be no doubt that it is an important factor in the movement of substances into and out of the cell. But the passage of many substances across the cell membrane in concentrations high enough to kill the protoplasm follows only upon the destruction of, or irreversible changes in the surface membrane and the consequent more or less complete disappearance of its special physiological characteristics. Furthermore, the cell membrane is alive and therefore metabolically active to some extent, and its peculiar properties as regards permeability are dependent upon the fact that it is alive, since they disappear with its death. It is evident then, first, that semi-permeability is more or less closely associated with metabolic conditions and second, that differential susceptibility to external agents cannot be interpreted in terms of permeability as distinct from metabolic conditions.

Experimental analysis has already shown that many of the so-called cases of specific action of external agents are not properly speaking specific effects at all, but actually are differences in degree rather than in kind. And the question may be raised whether at least many other apparently specific effects will not prove to be of the same sort.

Even admitting that different agents do attack the protoplasmic reaction system at different points—as they undoubtedly do—the important point is that in such a complex and inter-correlated physico-chemical reaction system as protoplasm, it is hardly possible to alter any one set of reactions or conditions to any considerable degree, without disturbing the system as a whole. Consequently in subjecting the frog egg or any other developing organism to an agent that in any way inhibits protoplasmic activity to a sufficient degree, it is certainly, in its broader aspects, not a specific, but a *general* or quantitative effect and reaction of the organism to the agent that becomes apparent. It is the striking uniformity of the susceptibility relations of organisms to so many different agents and conditions that forces the conclusion that there must be some fundamental feature of all axiate organisms that is affected in much the same way by different agents. In other words, *it is the existence of a gradient or gradients in metabolic rate, and protoplasmic conditions associated with them along physiological axes in the organism, that determine the uniformity and the susceptibility relations that have been shown to exist, and afford the basis for a rational interpretation of teratological development.*

VI. SUMMARY.

1. In the ovarian egg of the frog there is a definite relation between the polar axis of the egg and its blood supply. In every case the pigmented hemisphere is more richly supplied with arterial blood than the unpigmented hemisphere. It is suggested that polarity arises in response to this external factor, viz., respiratory and nutritive relation of the egg to the parent body. It also seems probable that the localization of the pigment on the egg is in response to the greater oxygen supply over a restricted area. There is no orientation of the ovarian egg to gravity.

It is pointed out that polarity in a number of other organisms arises in response to conditions external to the egg.

2. It is shown that differences in susceptibility exist in different parts of the egg and embryo of the frog. These differences are evidenced in the following ways:

(a) By differential death gradients,

(b) By differential inhibition, and

(c) By differential acclimation and recovery.

3. The two modifications of the susceptibility method described in this paper, (a) and (b) above, agree in showing that:

(a) Those parts of the egg where development first begins and proceeds most rapidly die soonest in lethal concentrations or intensities of external agents and are inhibited most under conditions so severe that acclimation does not occur. Both disintegration processes and inhibition are differential, paralleling the polar axis in early development and other axes that arise later.

(b) All of the modifications, produced under different conditions, are essentially similar, differing not in kind but in degree. No evidence was found of "specificity," "blastolysis" or "disorganization" as effective factors in the production of any of the modifications.

(c) The experimental data indicate that any type of abnormality may be produced under the influence of any inhibiting agent by controlling the concentration or intensity of action, the length of exposure and the stage in development (physiological condition) of the egg or embryo when exposed.

4. Differential inhibitions, appearing under conditions that prevent acclimation, are evidenced in the following and other ways:

(a) By the relatively slower division of the more active animal pole cells with the result that the cell size ratio approaches and may become equal to one.

(b) By the retardation or prevention of downward movement of material from the animal hemisphere, which takes place, normally, most rapidly in the sagittal plane, but which under these conditions, is most retarded in this plane.

(c) By the retardation of the dorsal lip region with the formation of V-shaped blastopores and oval yolk plugs that may be several times longer in the sagittal plane than broad.

(d) By the retardation of both dorsal and lateral lips of the blastopore resulting in the formation of flat-crescent blastopores and a whole series of modifications between this and,

(e) Perfect equatorial gastrulae.

(f) By the appearance of a secondary invagination apical to the original blastopore.

(g) By the shifting of the blastopore so that it may come to lie at or above the equator of the egg. (Accompanies *d*, *e*, *f*.)

(h) By the apparent obliteration of bilaterality, resulting in the formation of embryos radially symmetrical about the original polar axis.

(i) In the later embryonic stages a variety of abnormal types appear most of which are later stages of *c*, *d*, *e* and *g*, above. Among them may be mentioned, microcephalic forms with eyes, nasal pits, and ventral suckers in all degrees of approximation to complete "fusion"; anencephalic forms appear, and spina bifida of all degrees are common. Most of the forms show a marked dorsal concavity and the medullary folds may remain open in the brain region or throughout their entire length.

5. Since the dorsal lip region arises at the distal end of a rapidly growing region, and bears a definite spatial relation to the apical region that is experimentally controllable, it is probable that it arises by physiological isolation consequent upon the rapid growth in length, especially in the sagittal plane.

6. The data presented indicate that at the beginning of development, metabolic processes are most rapid at the apical (animal) pole and decrease toward the basal (vegetative) pole.

7. With advance in development a secondary region (dorsal lip) of high susceptibility appears, probably as a physiological isolation and represents a posterior "segmental" region of rapid growth similar in many respects to the posterior growing region of annelid larvæ.

8. It is pointed out that the existence of gradients in metabolic rate along physiological axes in the egg and embryo make possible the great uniformity and lack of specificity shown to exist in the susceptibility relations in the frog and other organisms, and affords the basis for a rational interpretation of teratogeny, not only in the frog but also in vertebrates generally.

VII. LITERATURE.

Bataillon, E.

'01 Etudes experimentales sur l'Evolution des Amphibiens. Arch. Entw.-Mech., 12: 610-655.

Behre, E. H.

'18 An experimental study of acclimation to temperature in *Planaria dorotocephala*. BIOL. BULL., 36: 277-317.

Child, C. M.

- '15a Individuality in organisms. Univ. of Chicago Press.
 '15b Senescence and rejuvenescence. Univ. of Chicago Press.
 '15c Axial gradients in the early development of the star fish. Amer. Jour. Physiol., 37: 203-219.
 '16a Axial susceptibility gradients in the early development of the sea-urchin. BIOL. BULL., 30: 391-404.
 '16b Experimental control and modification of larval development in the sea urchin in relation to the axial gradient. Jour. Morph., 28: 65-133.
 '16c Studies on the dynamics of morphogenesis and inheritance in experimental reproduction. IX. The control of head form and frequency in *Planaria* by means of potassium cyanide. Jour. Exp. Zoöl., 21: 101-126.
 '16d Axial susceptibility gradients in algæ. BIOL. BULL., 31: 419-440.
 '16e Axial susceptibility gradients in algæ. Bot. Gaz., 62: 89-114.
 '17a Differential susceptibility and differential inhibition in the development of polychete annelids. Jour. Morph., 30: 1-63.
 '17b Susceptibility gradients in the hairs of certain marine algæ. BIOL. BULL., 32: 75-92.
 '17c Gradients in susceptibility to cyanides in the meridional conducting path of the ctenophore *Mnemiopsis*. Amer. Jour. Physiol., 43: 87-112.
 '17d Experimental alteration of the axial gradient in the alga, *Griffithsia Bornetiana*. BIOL. BULL., 32: 213-233.
 '19a Demonstration of the axial gradients by means of potassium permanganate. BIOL. BULL., 36: 133-147.
 '19b The effect of cyanides on carbon dioxide production and on susceptibility to lack of oxygen in *Planaria dorotocephala*. Amer. Jour. Physiol., 48: 372-395.

Child, C. M., and Hyman, L. H.

- '19 The axial gradient in Hydrozoa. BIOL. BULL., 36: 183-221.

Gurwitsch, A.

- '95 Ueber die Einwirkung des Lithionchlorids auf die Entwicklung der Frosch- und Kröteneier. Anat. Anz., 11: 65-70.
 '96 Ueber die formative Wirkung des veränderten chemischen Mediums auf die embryonale Entwicklung. Archiv. Entw.-Mech., 3: 219-260.

Hargitt, C. W.

- '15 Regenerative potencies of dissociated cells of *Hydromedusa*. BIOL. BULL., 28: 370-384.

Hertwig, O.

- '92 Urmund und Spina bifida. Archiv. f. Mikrosk. Anat. 39: 353-503.
 '94 Ueber den Einfluss äusserer Bedingungen auf die Entwicklung des Froscheies. Sitzungsber. d. Akad. d. Wissen. zu Berlin.
 '95 Beiträge zur experimentellen Morphologie und Entwicklungsgeschichte (1) Die Entwicklung des Froscheies unter dem Einfluss schwächere und stärkerer Kochsalzlösungen. Archiv f. Mikrosk. Anat., 44: 285-344.
 '96 Handbuch der vergleichenden und experimentellen Entwicklungslehre der Wirbeltiere. Bd. 1, Pt. 1, 1st half, pp. 967-998.

Hyman, L. H.

- '16a An analysis of the process of regeneration in certain microdrilous ologochætes. Jour. Exp. Zoöl., 20: 99-163.
 '16b On the action of certain substances on oxygen consumption. I. The action of potassium cyanide. Amer. Jour. Physiol., 40: 238-248.

- '17 Metabolic gradients in *Ameba* and their relation to the mechanism of ameoid movement. *Jour. Exp. Zool.*, 24: 55-99.
- '19 On the action of certain substances on oxygen consumption. II., Action of potassium cyanide on *Planaria*. *Amer. Jour. Physiol.*, 48: 340-371.
- Jenkinson, J. W.**
- '06 On the effect of certain solutions upon the development of the Frog's egg. *Archiv Entw.-Mech.*, 21: 367-460.
- '09 Experimental embryology. Oxford.
- '11 On the effect of certain isotonic solutions on the development of the frog. A correction and extension of previous observations. *Archiv. Entw.-Mech.*, 32: 688-698.
- Kellicott, W. E.**
- '13 Chordate Development. Chapter II. Henry Holt and Co., N. Y.
- '16 The effects of low temperature upon the development of *Fundulus*. A contribution to the theory of teratogeny. *Amer. Jour. Anat.*, 20: 449-482.
- King, H. D.**
- '02 The gastrulation of the egg of *Bufo lentiginosus*. *Amer. Nat.*, 36: 527-548.
- Kopsch, F.**
- '00 Ueber das Verhältnis der embryonalen Axen zu den drei ersten Furchungsebenen beim Frosch. *Intern. Monatschr. Anat. Physiol.*, 17: 1-25.
- Lewis, W. H.**
- '09 The experimental production of cyclopia in the fish embryo (*Fundulus heteroclitus*). *Anat. Rec.*, 3: 175-182.
- McClendon, J. F.**
- '12 An attempt towards the physical chemistry of the production of one-eyed monstrosities. *Amer. Jour. Physiol.*, 29: 289-297.
- Morgan, T. H.**
- '97 The development of the Frog's egg. The Macmillan Co.
- '03 The relation between normal and abnormal development of the embryo of the frog as determined by the effect of lithium chloride in solution. *Archiv Entw.-Mech.*, 16: 691-712.
- '06a Experiments with Frog's eggs. *BIOL. BULL.*, 11: 71-92.
- '06b Origin of the organ-forming materials in the Frog's embryo. *BIOL. BULL.*, 11: 124-136.
- Morgan, T. H., and Boring, Alice M.**
- '03 The relation of the first plane of cleavage and the grey crescent to the median plane of the embryo of the Frog. *Archiv. Entw.-Mech.*, 16: 680-690.
- Morgan, T. H., and Stockard, C. R.**
- '07 The effects of salts and sugar solutions on the development of the Frog's egg. *BIOL. BULL.*, 13: 272-279.
- Morgan, T. H., and Tsuda, Ume.**
- '94 The orientation of the Frog's egg. *Quart. Jour. Micr. Soc.*, 35: 373-405.
- Newman, H. H.**
- '17 On the production of monsters by hybridization. *BIOL. BULL.*, 32: 306-321.
- Schultze, O.**
- '95 Ueber die Einwirkung niederer Temperatur auf die Entwicklung des Frosches. I. *Anat. Anz.*, 10:

- '99 Ueber die Einwirkung niederer Temperatur auf die Entwicklung des Frosches. II. Anat. Anz., 16.

Stockard, C. R.

- '09 The development of artificially produced cyclopean fish. "The magnesium embryo." Jour. Exp. Zoöl., 6.
- '10 The influence of alcohol and other anæsthetics on embryonic development. Amer. Jour. Anat., 10.
- '14 The artificial production of eye abnormalities in the chick embryo. Anat. Record, 8: 33-41.

Werber, E. I.

- '15 Experimental studies aiming at control of defective and monstrous development. A survey of recorded monstrosities with special reference to ophthalmic defects. Anat. Rec., 9: 529-562.
- '16 Experimental studies on the origin of monsters. I., An etiology and an analysis of the morphogenesis of monsters. Jour. Exp. Zoöl., 21: 485-584.
- '17 Experimental studies on the origin of monsters. II., Regarding the morphogenesis of duplicities. Jour. Exp. Zoöl., 24: 409-443.

Wilson, C. B.
















- '97 Experiments on the early development of the amphibian embryo under the influence of ringer and other salt solutions. Archiv. Entw.-Mech., 5: 615-648.
















Wilson, H. V.


- '07 On some phenomena of coalescence and regeneration in sponges. Jour. Exp. Zoöl., 5:
- '11 On the behavior of dissociated cells in Hydroids, *Alcyonaria* and *Asterias*. Jour. Exp. Zoöl., 11:

ERRATA.

Plates illustrating article by G. W. Tannreuther in the September issue. Should be inserted following page 208.







BIOLOGICAL BULLETIN

THE ORANGE STRIPED ANEMONE (*SAGARTIA LUCIÆ*, VERRILL). AN ECOLOGICAL STUDY.

LEON AUGUSTUS HAUSMAN, PH.D.,
CORNELL UNIVERSITY.

The orange striped anemone (*Sagartia luciæ* Verrill) is one of the commonest of the marine invertebrates to be found along the Connecticut coast. It was first observed at New Haven in 1892 by Miss Lucy Verrill, and described in 1898 under the name of *Sagartia luciæ* by Professor Verrill, who believed that it had been introduced into this locality from more southerly waters of North America, possibly on the shells of oysters. From New Haven it has greatly extended its range, and in 1901 was reported as being found as far north as Salem, Mass.

The morphology and anatomy of *Sagartia luciæ*, as well as its general physiology and geographical dispersal, have been ably dealt with in earlier papers. The present contribution embodies the results of a summer's observations made at various points along the Connecticut shore of Long Island Sound, and is chiefly concerned with the ecological relationships of the creature.

Although *Sagartia luciæ* is one of the most numerous of the invertebrates of the littoral fauna of Long Island Sound, it is, nevertheless, an inconspicuous member of the fauna, both because of its diminutive size (being normally about a quarter of an inch, or less, in diameter) and because of the concealing character of its olive green coloration. The bright orange stripes, to the number of about twelve in adults, further aid in rendering the creature inconspicuous by interrupting the contour of the body. The tentacles are usually arranged in four rather ill-defined rows, each row consisting of twelve tentacles. Within each row there is contained one tentacle, larger and longer than the rest, which normally corresponds in location with an orange

stripe. These larger tentacles, also, indicate the position, within the body, of the equally numerous intermesenterial spaces. Within the circle of tentacles lies the greenish oral disc, the center of which is pierced by the mouth opening, surrounded by the lip band. The latter structure is usually reddish in color; is impressed by a number of folds about its inner edge; and bears two prominent, opposite, gonidial grooves.

Since reproduction is often accomplished by longitudinal fission, or less commonly by budding, or by dividing at the base in a sort of budding process which Davenport (1) has termed basal fragmentation, individual anemones vary considerably in form and size, according to the stage of life which each is passing through at the time of the observation. Mature individuals which have not yet apparently begun reproductive activities, are illustrated by Figs. 1 and 2. Extremely unsymmetrical, and even monstrous, forms are frequently encountered, the results of accidental mutilation, for regeneration of lost parts reaches, with *Sagartia*, a high degree of development. Consequently surprisingly small fragments often settle down and regenerate an entire individual. Davenport has pointed out that new tissue, in individuals which are the results of reproduction by division, is of a hue of green less olive in tone than that of older individuals, and that the new stripes are a brighter yellow, with less orange tint. This is also true of new tissue which is being regenerated to make up for some lost portion of the body.

The general color scheme of *Sagartia* seems to be particularly well adapted to the concealment of the creature. The olive green of the body, broken by the longitudinal orange stripes, and the grayish white of the tentacles, forms a color aggregation of unusually subdued character when the animal is viewed in its normal habitat. When fully expanded, and hence in its most conspicuous attitude (Fig. 1), the body becomes lighter in color (because the more translucent) and the body assumes a translucency, as do also the tentacles. The latter, indeed, become, often, virtually colorless, and almost transparent. The color of the orange stripes likewise undergoes a diminution in intensity as the expansion of the animal progresses. Under these conditions, when viewed either from above or below (the latter by

means of a mirror placed at the bottom of the pool), the body outline becomes almost impossible to trace. Another aid to concealment is afforded by the immobility of the tentacles when the animal is fully expanded, and also by the extreme deliberation with which the act of expansion takes place. When contracted (Fig. 2) the animal appears like a small dark greenish button-like protuberance, whose outline, however, is interrupted by the orange stripes. This combination of dark olive green and orange (which seems at first a rather gaudy *ensemble*) can be shown to be a peculiarly effectual one for the type of rock on which *Sagartia* is usually found seated, *i.e.*, the granites of the New England coast. The rock is flecked with black, and with varying tints and shades of gray, green, reddish, and yellow, from the included quartz, feldspars, hornblende, etc., and from the various chemical combinations with these minerals which the oxygen of the air and the substances of the sea water make.

Sagartia luciae occurs below tide level on piles, or other submerged objects, and is frequently associated with the rock barnacle (*Balanus balanoides*), and with two species of mussels; the ribbed mussel (*Modiola plicatula*), and the black or edible mussel (*Mytilus edulis*). In the same company may also be found small clusters of the common oyster (*Ostrea virginica*). More often, however, the habitat of *Sagartia* are the tidal pools between the tide lines, among rocky headlands and shores, and rocky tidal islands, in which the New England coast abounds. Wherever it occurs, however, it will usually be found attached to vertical, or overhanging surfaces, rather than to horizontal ones (Fig. 3). In some few tidal pools it was found, however, on the gently sloping sides. Comparatively few enemies, either organic or inorganic, attack those individuals which are attached to the vertical or overhanging sides of the pool, while those that become detached and cast to the bottom are soon either buried by debris brought in by wave or tidal currents, or fall an easy prey to mud crabs (*Panopeus*), rock crabs (*Cancer irroratus*), or hermit crabs (*Eupagurus longicarpus*), for *Sagartia*, despite its armament of nettling acontia, is without means of defense against these heavily armored scavengers.

As a rule, the rock itself seems to be the favorite location of

Sagartia, for the creature can detach itself, at will, from its seat, and move by means of its tentacles, or float about, head downward, from the surface film of the water until a more favorable locality is found. The fact that the rock, or other seat of equal solidarity, is chosen more often than any other for the obsession of *Sagartia* would seem to indicate that the creature possesses a sense which instructs it regarding the relative stability of submerged objects. However it is often found attached to the shells of living mussels and even sometimes upon eel grass, fucus, or other algæ. It often occurs, also, attached to shells, and other objects that have become firmly wedged into crevices in the rock by the action of the waves.

Sagartia is ordinarily solitary in its habit, and not communal, and yet, what seem to be in effect, colonies, at least in so far as mere physical propinquity is concerned, were met with in tidal pools, where, apparently by repeated divisions, coupled with but little migration on the part of the increasingly numerous progeny, groups of from ten to twenty-five individuals, with their bases almost in contact, were formed. That this group formation aided in securing food was seen when large beach fleas (*Orchestia agilis*), which were easily able to escape the embrace of a single anemone, were at once, caught and held by the numerous tentacles brought to bear in the capture by several contiguous individuals at once, and easily nettled into a state of helplessness. Possibly such a colonial existence (if it may be so called) makes also for the more effectual protection of the individual members of such a community. Such groupings seemed to be the exceptions, and not the rule, in the disposition of the anemones in the tidal pools. It is interesting to conjecture, that possibly this grouping, perchance accidental at first, marks the beginning of the development of a movement toward communal living on the part of *Sagartia*. If this manner of living does make for the better protection of the members of colony, and aids in securing more frequent and larger captures of food, then we may suppose that the collective individual will thrive at the expense of the independent one, a law of progress derivable from other fields also than the biologic one.

The food of *Sagartia* consists mainly of small crustacea,

annelids, etc., which are seized by the tentacles, rendered innocuous by the acontia and then gradually manipulated into the mouth. Fig. 4 represents an individual capturing a beach flea, which had been steered into the blossom of tentacles with a broom splinter. All sorts of sufficiently soft food substances, either living or dead, are taken indiscriminately. Numerous individuals were fed by the writer, both in their natural habitats, undisturbed, and in aquaria, with pieces of fish, both fresh and decaying; clam, mussel, beef, bread, salt pork, insects, etc., which the anemones seized and devoured with the same impartial gusto which they showed for living creatures. They however rejected hard substances, such as sand grains, bits of shell, wood, etc. Beef juice, dexterously squirted upon one portion of the blossom of tentacles with a finely drawn out pipette, caused them to respond at once by throwing over a large number of tentacles to that side whence the stimulus had come and endeavoring to entrap some object. Frequently the tip of the pipette would be grasped, only to be relinquished again. Foreign substances, *i.e.*, those not usable as food, were pushed to one side of the blossom of tentacles and allowed to fall off. The food which seems to form the bulk of their intake consists of the smaller crustaceans, such as small beach fleas (*Orchestia agilis*), small clam worms (*Nereis*), and other small marine worms, very small crabs in the soft shelled stage just after a molt, chiefly the mud crab (*Panopeus*), crushed rock barnacles (*Balanus balanoides*), very small fish fry, and heterogeneous particles of animal tissue which the tidal and wave currents chance to float by. A lessening of the food supply in some particular neighborhood apparently is the moving cause that sets many individuals roving by hanging head downwards from the surface film of the water and being borne here and there by currents until a more fruitful locality is discovered. Rarely there was found a more ambitious individual which had fastened upon the shell of a periwinkle (*Littorina littoria*), and was continually being carried into pastures new.

The enemies of *Sagartia* are numerous. Among them the most important are: various species of rock bottom feeding fishes, rock and mud crabs, starfishes (*Asterias*, and others), and the

larger clam worms. Many individuals are doubtless swept away by the assaults of sand and debris-laden waves in times of storm, and by the grating of ice cakes against the rocks during the winter. Frequently a tidal pool is drained by an excessively low ebb tide, and the anemones left exposed for an hour or so to the rays of a direct sun. This, coupled with a vigorous dry wind will often accomplish the destruction of some individuals, though they were able to withstand even such adverse conditions to a surprising degree, as experiments showed.

It seems remarkable that so soft and relatively defenseless a creature should have so greatly increased its numbers and extended its range in our waters. Its success in meeting the unusually numerous vicissitudes of a littoral existence may perhaps be attributed to the following: (1) in ability to withstand considerable differences in temperature, (2) its ability to withstand buffeting by the waves because of the yielding and resilient character of its body, (3) its ability to contract tightly, and to survive through a period of foul water, or of dry conditions exposed to the sun and wind, (4) its apparent disregard of differences in the salinity of the water, (5) its protective coloration, (6) its defensive acontia, (7) its rapid rate of reproduction and growth to maturity, (8) its several methods of reproduction, (9) its ability to withstand annihilation through laceration, and (10) its ability to regenerate lost parts.

BIBLIOGRAPHY.

1. Davenport, A. G.
'03 Variation in the Number of Stripes of the Sea Anemone (*Sagartia luciae*). Mark Anniv. Volume, p. 137.
2. Carlgren, O.
'93 Studien über Nordische Actinien. Kongl. Svenska Vet. Akad. Hdlgr. N. F., Bd. 25, No. 10, p. 1.
3. Dixon, F.
'88 On the Arrangement of the Mesenteries in the genus *Sagartia* Gosse. Sci. Pro. Roy. Dublin Soc., n. s., Vol. 6, p. 136.
4. Dixon, G. Y. and A. F.
'91 Report on the Marine Invertebrate Fauna Near Dublin. Pro. Roy. Irish Acad., S. 3, Vol. 2, p. 19.
5. Gosse, P. H.
'60 Actinologia Britannica, Lond. 8mo, pp. xl + 362, Pls. 12.
6. Lang, A.
'86 *Gastroblasta Raffaelei*. Jena. Zeit. für Naturwiss., Bd. 19, S. 20.

7. McCrady, J.
 - '59 Instance of Incomplete Longitudinal Fission in *Actinia cavernosa*. Bosc. Pro. Elliott Soc. Nat. Hist., Charleston, S. C., Vol. 1, p. 275.
8. McMurrich, J. P.
 - '91 Contributions on the Morphology of the Actinozoa. III. Phylogeny of the Actinozoa. Jour. Morphol., Vol. 5, No. 1, p. 125.
9. Parker, G. H.
 - '02 Notes on the Dispersion of *Sagartia lucia*. Am. Nat., Vol. 36, p. 491.
10. Verrill, A. E.
 - '98 Description of New American Actinians, with Critical Notes on Other Species. I. Am. Jour. Sci., Ser. 4, Vol. 6, p. 493.

EXPLANATION OF PLATE I.

FIG. 1. A fully expanded, mature *Sagartia* with its tentacles in exploring posture, after having been stimulated with beef juice injected into the water. The object to which it is attached is an edible or black mussel (*Mytilus edulis*).





FIG. 2. Contracted mature *Sagartia*.

FIG. 3. Transection through a typical ideal tidal pool, to show the disposition of *Sagartia* individuals, and their general ecological relationships.

1. *Sagartia lucifera*.
2. Edible or Black Mussel (*Mytilus edulis*).
3. Ribbed Mussel (*Modiola plicatula*).
4. Mud Flat or Elephant Snail (*Nassa obsoleta*).
5. Oyster Drill (*Urosalpinx cinerea*).
6. Periwinkle (*Littorina littoria*), not numbered in the figure. It occurs in great numbers, everywhere.
7. Oyster (*Ostrea virginica*) rare in tidal pools.
8. Clam Worms (*Nereis*), in mud among masses of mussels.
9. Small Ribbon Worms (*Mechelia*).
10. Starfish (*Asterias forbesi* and *vulgaris*).
11. Hermit Crabs (*Eupagurus longicarpus*).
12. Mud Crabs (*Panopeus herbstii* and others).
13. Rock Barnacle (*Balanus balanoides*).
14. Green Beach Flea (*Orchestia agilis*).

FIG. 4. *Sagartia*, capturing a small Beach Flea (*Orchestia agilis*).

FIG. 5. Partially contracted *Sagartia*, of large size, just before fission. A strong light had been placed behind the creature, revealing the manner of the accommodation of the contracted tentacle blossom, and the folded sides of the gastrovascular cavity. The latter is seen to be empty of food.



PRELIMINARY NOTE ON THE SPERMATOGENESIS OF *PEDICULUS VESTIMENTI*.¹

KATHARINE FOOT.

In response to a request from Col. Alexander Lambert I came to Paris, December, 1917, in order to study *Pediculus vestimenti* from the point of view of the biologist in the hope of getting some data that might be of service to the investigators who were studying the louse as a possible transmitter of certain diseases prevalent among the troops.

In an exhaustive and masterly study of the problem in relation to trench fever, Col. Strong has shown that the lice can unquestionably transmit this disease and has proved beyond question what was heretofore merely a surmise.

Such data as I have been able to collect that may have some bearing on the problems in relation to disease have been reported to the Research Department of the American Red Cross; but I have omitted from such reports purely cytological data which have no apparent practical value.

As far as I know, no report has been published of any work done on the chromosomes of the louse, and as *P. vestimenti* has been classed as an Hemipterous insect I was interested to see if the chromosomes have the same bizarre morphological characteristics as are typical of so many Hemiptera.

The spermatocyte chromosomes are very minute, so minute that I have as yet found it impossible to demonstrate the method of division of the second spermatocytes; but I have several entirely satisfactory stages of the first spermatocyte chromosomes, and as these have the same morphological characteristics as the corresponding stages in other species of Hemiptera, it is quite permissible to assume a like correspondence for the second division.

¹ My grateful acknowledgments are due to the late Professor Blanchard, who gave me a most cordial welcome to his laboratory at the Ecole de Médecine. He not only gave me ample space for my work but the sympathy and encouragement I received from him and the members of his staff made it possible for me to continue my investigations of these repulsive insects.

The above mentioned morphological peculiarity of the chromosomes of many Hemiptera *i. e.* the unequal bivalent—called by Wilson the X Y chromosome—is present in the first spermatocytes of *Pediculus vestimenti* and its division is typical—the two unequal parts dividing as univalents in the first division. The unequal bivalent is demonstrated in the metaphase stages of Figs. 10, 11, 12, and the division of the smaller half of the unequal bivalent is shown in Fig. 13.

Miss Strobell and I (1914) published a series of photographs showing both the first and second spermatocyte divisions in *Euschistus variolarius*, *Euschistus servus* and in two generations of hybrids from *E. variolarius* by *E. servus*. A comparison of these photographs of the first spermatocyte chromosomes with the above mentioned Figs. 10–13 will show that the first spermatocyte chromosomes of *Pediculus vestimenti* are of the same type as in the species of *Euschistus* referred to, and it is therefore quite logical to assume that the second spermatocyte chromosomes are equally typical though its demonstration is not yet possible.

In Fig. 16 (on the right) is a group of chromosomes from an embryo in an egg at the basal end of the ovary.

It is not possible to determine the exact number of chromosomes of this group, as 18, 19 or 20 can be counted.

The first spermatocyte chromosomes indicate that the somatic number should be ten and in the few spermatogonial groups I have found it possible to identify ten chromosomes; but they are so small and so frequently constricted that the estimate can always be questioned.

I am convinced that the most favorable stage for an exact interpretation of the louse chromosomes is the first oöcyte prophase, but to secure this stage involves a patient search which cannot be undertaken at present. Fig. 18 indicates that it will be possible to find the later prophase stages, for in this preparation the chromosomes are nearly formed, and it ought to be possible to find the slightly later stages in the same locality of the ovary.

An early stage of the yolk spheres of the ovary is shown in the photomicrograph of Fig. 16 (on the left). These are chromosome-like structures, which in the early stages select the chroma-

tin stains and are morphologically so like chromosomes that one is tempted to interpret them as chromosomes which develop into yolk-spheres.

A series of stages prior to the first spermatocyte metaphase is shown in Figs. 1 to 9. The centrosomes of the first spermatocytes (Figs. 1 to 4) are a feature of these stages which is most marked. The position of these centrosomes in relation to the nucleus is very variable: they are in contact with the nucleus or almost in contact with the periphery of the cell or in any position between these two extremes. In later stages they are frequently divided into two or more, rarely into three, parts. Several of the former are shown in Fig. 7 (the centrosomes of this figure are a little exaggerated by the artist).

The Development of the Spermatid into the Mature Spermatozoön.—The most striking feature in the development of the spermatozoön is the duplex character of the tail. Apparently the tail is composed of two distinct and independent filaments, the apparent independence of these filaments being more marked in the earlier stage (Figs. 20–21). One would naturally suppose that one of these filaments represents the flagellum of other forms, but it has not been possible to demonstrate any substance connecting the two filaments, though the fact that even in smear preparations the two are never found widely separated would indicate that they are attached by some connecting substance.

HISTORICAL SKETCH.

Certainly no insect has been accused of being the promoter of a greater variety of diseases than the louse and perhaps no accused has been charged with so many crimes on less evidence. *Pediculus* has been credited with transmitting the following diseases: typhus fever, typhoid fever, recurrent fever, trench fever, tuberculosis, spinal meningitis, plague, leprosy, beri-beri and more than a dozen minor skin diseases. In some of these cases the evidence seems to be confined to the fact that the patient may be infested with lice.

For typhus fever, recurrent fever and trench fever it has been proved that the lice do in fact transmit these diseases, but details as to the method of transmission are still disputed. It is held

by many investigators that germs are not transmitted by the bite of the louse—the sole method of transmission being infection from their excrements. These are freely deposited on the skin and in the clothing of the host and subsequent scratching of the skin induced by the intense itching of the bites not only lacerates the surface but frequently causes a deep wound that leaves a scar lasting many months. It is self-evident that such lacerations over surfaces more or less infested with the excrements of the lice may cause a most effective inoculation. This method of infection by the faeces or the crushed body of the louse has been demonstrated for typhus fever by Nicolle (1909), Nicolle, Conte and Conseil (1910) and others. For recurrent fever by Sergent and Foley (1910), Sergent, Gillot and Foley (1911), Nicolle, Blaizot and Conseil (1912) and others. For trench fever, by R. P. Strong (1918). Familiarity with the feeding habits of the louse demonstrates the danger of this method of inoculation, for the amount of excrement discharged by each louse is surprising. I have frequently seen a single louse, during one hour's feeding, discharge excrement ten times, and five times is not unusual. For more than a year I have closely studied the feeding habits of *Pediculus vestimenti* and in my report to the research department of the Red Cross I described their behavior as follows: "Observations made during the feeding hour demonstrate that individual lice may behave very differently. As a rule they bite at once when young and vigorous. Some become gorged with blood in ten minutes and will not bite again, though most frequently they bite several times during the hour, moving around rather restlessly between times. Others bite continuously the entire hour, casting their excrements while biting. The old lice frequently do not bite for several minutes or even half an hour and then suck the blood very deliberately." I am inclined to believe that the method of biting demonstrated for one hour indicates the method for the entire twenty-four hours and that therefore the younger lice are almost continuously feeding on the host, wandering about and biting very frequently.¹ This would accord with observations made by Miss Strobell and

¹ These observations support Nuttall's (1917) conclusions as to the probable feeding habits of lice. He thinks they bite very frequently, for when raising them on his wrist he noticed they started to bite at all times when he was quiet.

myself on other species of Hemiptera, *Euschistus variolarius*, *E. servus*, *E. ictericus*, etc., etc. These insects were fed on fruit and could be closely observed during the twenty-four hours. They fed almost continuously during the night as well as during the day, and this leads me to surmise that the lice may feed with equal frequency and explains the torment that soldiers suffer even when infested with relatively few lice and suggests the possibility of inoculation from a single louse.

The disputed question whether simply the bite of the louse can inoculate the host has given rise to much discussion, some investigators emphatically denying that any danger is caused by the bite alone. Colonel Strong (1918) conducted some experiments with the aim to determine this point and concludes that "it seems fair to argue that the bite is probably a common mode of infection." He states that in some instances the disease was produced by pure biting experiments. His summary of the probable methods of infection through biting is as follows:

1. By piercing or stabbing and inoculating with mouth-parts contaminated with infected material such as blood from the patient or by louse feces and body juices.
2. By stabbing and inoculating from the skin which has been contaminated with infected material such as louse feces, and possibly body juices.
3. By stabbing and inoculating with mouth-parts which have been contaminated with virus grown or developed in the stabber-sac.
4. By stabbing and regurgitating of the virus from the alimentary canal.
5. By stabbing and the injection of the virus contained in the salivary juices.
6. By hereditary infection.

One of the difficulties in determining the value of the bite alone is to eliminate the feces from the experiment. Those cases in which this is assumed to be done by allowing the louse to bite through chiffon do not appear to me to be conclusive, for in my experience they will not bite unless the chiffon is pressed upon the surface with sufficient strength to force the skin through the interstices of the chiffon, in which case the

only effect of the chiffon is to reduce the area of the skin exposed, and the lice wander over the exposed area distributing the fæces as usual.

The danger of the louse as a promoter of disease has been so long appreciated that he has claimed the attention of a large number of investigators, the French and English forming the majority. The work accomplished up to 1917 has been most ably presented by Professor Nuttall, of Cambridge. His bibliographical list is an index of the thoroughness of his historical study of the subject. He has listed nearly 600 investigators.

A second historical sketch was published in France the same year (Souéges et du Noyer, 1917). These two studies are a convenient record of all the historical data that can be of value to the investigator.

Two English investigators (Warburton, 1909, and Fantham, 1912) were the first to study the life history of *Pediculus* and their results were supported and extended by Bacot in 1916. He determined the number of moults to be three, the average length of life of the louse, the average number of eggs deposited daily by a single female and other details, all of which my investigations support although our methods of work differed materially. He used an entomological box containing a number of lice and strapped this box on his person each night, allowing the lice to bite from six to seven hours daily. My lice were fed only one hour in the twenty-four and in such a manner that I could watch them while feeding.

The most serious difficulty in the investigation of lice is the food supply. In all the accounts with which I am familiar the investigator has had sufficient self-abnegation to feed his lice on his own person, but not having reached those heights myself, my initial difficulty was to find a host. There seems to be something extremely ridiculous in the mere suggestion of feeding a louse, for my most serious and generous offers received the discouraging response of a broad grin and an emphatic shake of the head. I finally succeeded in securing a host at the Asile de Nuit—a night employee of that institution. He was an old sailor whose evident familiarity with *Pediculus* at the Asile de Nuit had led him to cease to regard them in a humorous light,

and he proved to be a thoroughly dependable food supply. He never missed his daily hour in my laboratory for the five months I employed him.

When feeding the lice I at first used the usual method of putting a number in a tube, inverting the tube on the arm and holding it securely in place to prevent the lice from escaping. I found this method unsatisfactory for several reasons and devised therefore quite a different technique. Lice cannot crawl up a glass surface if it is clean and are therefore perfectly safe in a glass ring even if it is only 2 cm. high. I had such rings made to order and fastened them securely onto the arm with melted paraffine. In this manner several different experiments can be conducted at the same time and the generations can be kept separate—further the lice can be conveniently studied with a lens during the hour they are feeding. For the remaining twenty-three hours they were kept in a Pasteur incubator at a temperature between 27° and 29° C. While in the incubator the lice were kept in cages such as those used in the laboratory for raising various insects. This cage is the tube de Borel, in which is placed an inner tube for the insects, this being held in the center by absorbent cotton which is kept wet to insure sufficient moisture. I found the use of absorbent cotton very inconvenient and replaced it with a short tube having an aperture large enough to contain the inner tube and open at both ends with a lip at each end sufficiently wide to center it in the tube de Borel. The inner tube in which the insects are kept is dropped into this shorter tube and an inch of water kept in the tube de Borel. I found this method a great economy of time, for it was necessary to pack the cotton around the inner tube with much care, since if the opening came in contact with the inner surface of the tube de Borel (often quite wet) a drop or two sometimes dripped into the inner tube and cost the life of one or more nymphs.

Several years of experience in crossing and raising other species of Hemiptera have been my guide in raising the lice. Miss Strobell and I found that the species we studied required as much humidity as possible while avoiding any condensation of the moisture. This I have found true for lice—a half a drop of water or less can kill a nymph. If he gets on his back on the

glass in even a fraction of a drop of water, he cannot regain his feet until the water dries and if the glass is not clean he adheres to it and finally dies.

After trying the usual method of keeping the lice on small pieces of woollen or muslin cloth it occurred to me that a large number of short pieces (about 8 mm.) of soft, coarse thread would have many advantages. First they would be much more sanitary for they can be changed every day if necessary without disturbing the lice at all. When lice are on a small piece of cloth, the cloth becomes filthy in a few days and it is exceedingly difficult to remove the lice to fresh pieces. Further the thread avoids all the difficulties encountered in transferring and counting the lice. They cling to a thread with great tenacity; therefore single lice can be carried on a thread any distance with perfect safety. They deposit their eggs on the thread and therefore the eggs deposited each day can be conveniently collected and isolated. Using these threads made it a simple matter to record the following life history of a single pair of lice. The pair was hatched from eggs deposited in the laboratory and had their third (final) moult August 19. They were seen mating August 22. The next day 4 eggs were deposited and thereafter 4, 5 or 6 were deposited daily until the female died.

RECORD FL. D.

1918.		Eggs.	Hatched.
Aug.	(Mated Aug. 22)		
23	Mated.....	4	
24	3	
25	5	
26	4	
27	4	
28	5	
29	5	
30	4	
31	5	
Sept.			
1	5	
2	Hatched the 10th day.....	6	1
3	5	3
4	5	4
5	(Mated).....	5	3
6	(Male dead—each daily deposition of eggs kept separate from death of male).....	5	7
7	5	3

As the female had her third moult between the 18th and 19th of August, she lived only 28 days after maturity, less than the average length of life for a female; but other females in the laboratory kept under the same conditions lived 42 days, 40 days, 39 days, etc. Bacot found the average length of life of a female to be 34 days. If my lice are a little below this average I think it is probably due to the difference in feeding—he fed his lice six or seven hours daily and mine were fed only one hour daily.

According to my experience, the longer a race is bred in the laboratory the less prolific they become and the death rate is much higher. I believe this is due entirely to an abnormal lack of nourishment. Feeding only one hour in twenty-four is certainly very abnormal for these insects.

One cannot study the record of the large amount of experimental work done on the louse without being impressed with the need of feeding these insects apart from the human host before certain problems now in dispute can be solved. My efforts have been largely given to this well high hopeless task which is my apology for a very superficial study of the spermatogenesis.

PAPERS REFERRED TO.

Bacot, A.

'16 Notes on *Pediculus humanus* and *Pediculus capitis*. Brit. Med. Journ.

Foot, Katharine and E. C. Strobell.

'14 The Chromosomes of *Euschistus variolarius*, *Euschistus servus* and the Hybrids of the F₁ and F₂ generations. Arch. für. Zellf., Bd. 12.

Langeron, M.

'16 Précis de Microscopie. Masson & Cie, Editeurs, Paris.

Nicolle, C.

'09 Reproduction expérimentale du typhus exanthématique chez le singe. C. R. Acad. des Sc., 149.

Nicolle, C., Conte and Conseil, E.

'10 Transmission expérimentale du typhus exanthématique par le pou du corps. Ann. Inst. Past., 24.

Nicolle, C., Blaizot, L. et Conseil, E.

'12 Etiologie de la fièvre récurrente. Son mode de transmission par le pou. C. R. Acad. des Sc., 154.

Nuttall, George.

'17 Bibliography of *Pediculus* and *Phthirus*. The Part played by *Pediculus humanus* in the Causation of Disease. Parasitology, Vol. X.

Sergent, E. and Foley, H.

'10 Recherche sur la fièvre récurrente et son mode de transmission dans une épidémie algérienne. Ann. Inst. Past., 24.

Sergent, E., Gillot, V., et Foley, H.

'11 Typhus récurrent Algérien. Sa transmission par les poux. C. R. Soc. Biol., 70.

Soueges, R. et Du Noyer, M. R.

'17 Les Poux, le mal qu'ils nous causent. Bull. des Sciences Pharmacologiques t. XXIV.

Strong, Richard P.

'18 Trench Fever. Report of Commission Medical Research Committee. American Red Cross. Oxford University Press.

EXPLANATION OF PLATES.

All the sketches were drawn at a magnification of about 750. Zeiss hom. immer. 2 mm. 140—apo. oc. IV camera lucida.

TECHNIQUE.

Smear preparations stained with May-Grünwald followed by Hollande.

These stains were used in accordance with the method published by Dr. Langeron (1916) and I am greatly indebted to him for many valuable suggestions and encouraging interest in my work.

PLATE I.

FIGS. 1, 2, 3, 4. First spermatocyte rest stages, each showing a distinct centrosome in varying proximity to the nucleus. No cell membranes are differentiated.

FIG. 5. First spermatocyte nucleus showing a single nucleolus and the chromatin segregating to form the chromosomes.

FIG. 6. Two first spermatocyte nuclei with granular chromatin segregating to form the chromosomes. The nucleolus has disappeared.

FIG. 7. Numerous first spermatocyte nuclei showing successive stages of the differentiation of the chromatin. In the earlier stages the chromatin is apparently homogeneous and later it is granular and segregating into definite masses to form the chromosomes. A few centrosomes are differentiated.

FIG. 8. Numerous first spermatocyte nuclei showing later stages than those of Fig. 7. In many of the nuclei the chromatin has segregated into 5 distinct masses foreshadowing the 5 bivalent chromosomes of the first spermatocyte metaphase.

FIG. 9. First spermatocyte nuclei about the same stage of development as those of Fig. 7.

FIGS. 10, 11, 12 and 13. Each figure shows the 5 bivalent chromosomes of the first metaphase. In each figure one or more of the chromosomes is a dyad, foreshadowing the first division. An unequal bivalent, which is typical of so many Hemiptera, is clearly shown in each group.


FIG. 13. All the chromosomes of this first metaphase group are dyads, foreshadowing the division of each. The large and small chromosomes of the unequal bivalent are detached and each is a dyad. This indicates that each will divide in the first division and that therefore the resulting halves will undoubtedly separate in the second division in the manner typical of so many Hemiptera.

FIG. 14. Late anaphase of the second division. The chromosomes are too small and too closely segregated to determine their number and form.

FIG. 15. Three telophases of the second division. Each shows an unequal division of the chromatin, this being the sole evidence, at this stage, of the separation of the large and small moieties of the unequal bivalent.

FIG. 16. On the left a photomicrograph of a small group of chromosome-like structures from an immature ovary. From these the yolk-spheres are developed. $\times 450$. On the right a sketch of a group of chromosomes from an embryo in an egg at the basal end of the ovary.


FIGS. 17, 18. Two germinal vesicles from young ovarian eggs. In Fig. 17 the chromatin has partly segregated into threads, and a single nucleolus is present. Fig. 18 shows numerous small dense nucleoli, and the separate chromatin threads suggest a progressive step in the forming of the chromosomes though their abnormal number may be due in part to the technique.



EXPLANATION OF PLATE II.

FIG. 19. Numerous heads of spermatids shortly after the second division. They are at first round, as demonstrated in the figure, and, as a rule, elongate before the appearance of the tail. In the elongated heads of this figure the middlepiece (?) is in evidence and from this point the tail develops.

FIGS. 20 to 27. Successive stages of the development of the spermatid into the mature spermatozoön. In Figs. 20 and 21 the middlepiece is demonstrated, the tail developing from this point. In Figs. 22 and 24 both the middlepiece and the acrosome (?) are demonstrated, the spine developing from the latter. Fig. 25 shows an early stage of the development of the spine. Figs. 26 and 27 demonstrate the mature spermatozoön in which the head and spine are fully developed. At this stage the middlepiece and acrosome are obscure.



DETERMINATION OF THE SEX OF THE OFFSPRING FROM A SINGLE PAIR OF *PEDICULUS* *VESTIMENTI*.

KATHARINE FOOT.

Nuttall (1917) gives a summary of the data as to the proportion of the sexes in *Pediculus vestimenti*. He says "the proportion of the sexes as determined by raising experiments has yielded contradictory results," and this he thinks is due to the small number of the broods of the experiments. In his own experiments of mixed pairs he found the sexes nearly equal, though his broods also were small.

Hindle (1917) discovered a marked inequality in the proportion of the sexes of certain pairs, and he concludes this to be the normal condition, basing this conclusion on the determination of the sex of 25 single pairs.

It has been proved by the observations of several investigators that a single female may deposit from one hundred to nearly two hundred eggs. It is obvious, therefore, that the sex must be determined for at least one hundred of a generation or the results are inconclusive. Certainly at least 75 per cent. of the eggs deposited must hatch and the sex be determined for these nymphs.

Hindle secured such a small number of individuals from each of his 25 pairs that Nuttall seems more than justified in questioning his conclusions. The number of individuals in each of the 25 broods studied by Hindle is as follows: 2, 4, 6, 9, 8, 10, 11, 12, 15, 17, 24, 25, 26, 26, 27, 29, 30, 31, 32, 34, 36, 38, 45, 48, 64. Fifteen of these 25 broods were all males or all females, *e.g.*, the brood of two were both females, the brood of four were also females, one of the broods of nine were males and the other brood of nine were both males and females. The brood of sixty-four were all females but as they probably do not represent more than 50 per cent. of the offspring from that pair, the evidence, even in this case, is without force.

METHODS AND RESULTS.

In order to determine the sex of as large a number of the offspring as possible I adopted a different method from that of other investigators. Instead of waiting for the nymphs to mature in order to determine the sex, they were dissected at any stage that was convenient and those that died were not discarded but dissected at once and the sex recorded. It was thus possible to determine the sex of nearly all that were hatched.

Our method of treating these insects makes it possible to collect and isolate the eggs deposited each day, thus keeping an accurate count of the number of eggs deposited daily and the number hatched.

Both the male and the female had the third moult in the laboratory, the female July 22, 1919, and the male July 24.

The male died August 26; he lived therefore 33 days after maturity, thus exceeding the average length of life of *P. vestimenti* males. The female died September 5. She lived therefore 45 days after maturity, eleven days longer than the average length of life of the females. She continued to deposit eggs daily after the death of the male. She deposited in all 143 eggs (116 before the death of the male and 27 after that date). She commenced to lay eggs the 4th day after maturity and stopped laying 2 days before she died. She deposited 2, 3, 4 or 5 eggs each day. Of the 116 eggs laid before the male died, 11 failed to hatch, but nearly all developed and three of the nymphs partly emerged. Of the 27 eggs laid after the male died, seven failed to hatch, but all were fertilized and four of the nymphs partly emerged.

The nymphs were raised in lots of 20, and the death rate was relatively low in lots 1, 2, 3 and 4. The death rate in lots 5 and 6 was much higher. Several of the nymphs died even before the first moult. In some of these cases the nymphs were too dry to dissect, and therefore the sex was undetermined. This higher death rate was probably due to the age of the female, as the death rate was progressively high towards the end of the experiment.

125 eggs were hatched and the sex was determined for 115 (62 males and 53 females). 42 died during the experiment and

these were dissected as stated above and their sex determined. Of these 27 were males and 15 females.

For the first half of the experiment the females were more numerous than the males; the proportion of males then gradually increased until the final result gave 62 males and 53 females.

These results are of value only because the sex was determined for 92 per cent. of the entire generation, but the experiment must be repeated many times before the evidence can be of scientific value.

PHYSIOLOGICAL STUDIES ON PLANARIA.

III. OXYGEN CONSUMPTION IN RELATION TO AGE (SIZE) DIFFERENCES.

L. H. HYMAN,

HULL ZOÖLOGICAL LABORATORY, UNIVERSITY OF CHICAGO.

I. REVIEW OF LITERATURE.

The present paper is a report of some determinations of the rate of oxygen consumption of small (young) as compared with large (old) planarians belonging to three different species. That the metabolic and other activities of small individuals are more rapid than those of large individuals of the same and different species is so well known from a number of independent investigations that further evidence appears almost superfluous. The majority of these investigations, however, deal with vertebrates, where a certain difficulty inherent in the nature of the material is encountered. This difficulty centers about the impossibility of finding a proper unit for comparison of animals of different sizes. Among the warm-blooded vertebrates, the metabolic rate has commonly been reckoned per unit of surface area on the necessity of taking into account loss of heat by radiation. But leaving out of consideration the difficulties of accurately measuring the surface, over which proceeding a controversy of long standing exists, the surface method of comparison is more or less meaningless in the case of cold-blooded animals. Nor is the method of comparison of metabolic rate per unit weight free from objection. A considerable part of the weight of the vertebrates and the higher invertebrates consists of skeletal material, connective tissue, fat, etc. The metabolic rate of such material must be relatively low, and further its proportion to the total weight must be different in animals of different sizes; nevertheless, these factors have not been considered in any of the researches on the subject, nor, as far as I am aware, has anyone

attempted to make a correction for these inert portions of the body.

As Benedict and Talbot ('14) concluded after a study of the metabolism of infants, metabolism cannot be accurately measured either by weight or by surface, but only by the amount of active protoplasm. It is probably impossible to determine this but certainly in the soft-bodied lower invertebrates, weight is a more accurate index of the amount of active protoplasm than is the case in higher forms, where skeleton is present. For this reason, the metabolism of the lower forms deserves more attention than it has hitherto received from physiologists.

Among the early investigators of the effect of size differences on metabolic rate were Jolyet and Regnard ('77), who determined the rate of oxygen consumption per unit weight of a large number of fish, and some of the higher invertebrates. They found that among related groups small species consume more oxygen per unit weight per unit time than large species; and small individuals more than large individuals of the same species. Vernon ('95) confirmed this conclusion. He measured the rate of oxygen consumption per unit weight of large and small individuals of the same species, using as material a hydrozoan medusa, a scyphozoan medusa, two ctenophores, two gasteropods, and two pelagic tunicates. With a few exceptions the smaller individuals were found to consume relatively more oxygen than the large ones. Bounhiol ('02) working with twelve species of polychæte annelids, belonging to several different families, found that the carbon dioxide production was greater the smaller the individual. This was true for individuals of the same and of different species. On the other hand, Montuori ('13) was unable to come to any definite conclusion from his measurements of the oxygen consumption of a large number of species belonging to most of the aquatic groups. In some cases the small individuals respired relatively faster than large ones of the same species; in others the reverse was found; and in still others there was no relation between size and rate of respiration. The great irregularity of Montuori's results, as well as their disagreement with the work of others, suggests that he failed to control adequately the conditions under which the experiments were performed. Child

('19) found that small individuals of *Planaria dorotocephala* produce more carbon dioxide per unit weight than large ones, and this experiment is now regularly performed as a class experiment. Allen ('19) has shown that the rate of oxygen consumption of *Planaria agilis* per unit weight is greater the smaller the animals, and reports that Miss Wolf working in the same laboratory found the same to be true for the leech, crayfish, branchipus, may-fly nymph, and stone-fly nymph. It is also the case with dragon-fly nymphs, as shown by Mr. G. C. Hawk in this laboratory. According to Morgulis ('15), the oxygen consumed per gram per hour by flounders is in general greater for small than for large animals. Tashiro and Adams ('14) note that the carbon-dioxide production of the ganglionic cord of the heart of *Limulus* is relatively greater in small than in large cords. Nicolas ('18) found that the young leaves and stems of plants give off from $3\frac{1}{2}$ to 7 times as much carbon-dioxide per gram per hour as old leaves and stems from the same branch.

A considerable amount of labor has been devoted to this problem in the case of mammals, especially man. Rubner originally maintained that warm-blooded animals of different sizes produce the same amount of heat per unit surface, but subsequent investigations have shown that this point of view is erroneous. Thus Magnus-Levy and Falk ('99) clearly showed that the oxygen consumption and carbon-dioxide production is highest in children and decreases with age, as measured either per unit weight or per unit surface. If individuals of the same size and weight but of different ages are compared, the younger ones respire the faster. The authors conclude that the protoplasm of children has a definitely higher rate of respiration per unit weight than that of mature individuals. References to other work supporting these conclusions will be found in their paper. Recently Du Bois ('16) and Gephart and Du Bois ('16) have verified this earlier work. The graphs and tables presented by Du Bois show that the heat production as measured per kilogram of weight is highest in infants and decreases gradually; as measured per unit of surface, it is low in infants, increases rapidly during the first year, reaches a maximum between the ages of 1 and 6, falls rapidly to 20, and thereafter

decreases more slowly. According to Benedict and Talbot ('14, '15), the metabolism of infants is about the same as that of adults, but the results were very variable. It seems highly probable that the metabolism of infants cannot properly be compared with that at other ages because: (1), the heat-regulating mechanisms of infants are known to be very imperfect; (2), infants commonly have relatively more fat than is present in normal individuals of other ages; and (3), the muscle tone of the voluntary muscles of infants must be lower than it is at other ages. Since a large part of the heat production of mammals originates in the voluntary muscles, this difference in muscle tone alone makes impossible any real comparison of the metabolism of infants with that of later stages of ontogeny when the muscles are in full use.

A few researches have been carried out on mammals other than man. Thus Slowtsoff ('03) working on dogs found that the oxygen consumption and carbon dioxide production per kilogram per minute is greater in small than in large individuals. A. V. and A. M. Hill ('13) determined the same relation in rats. Among birds, Bohr and Hasselbalch ('00) observed that the carbon dioxide production per kilogram per hour is considerably higher in newly hatched chicks than in the adult hen.

These researches are sufficiently numerous to establish the generalization that smaller (younger) individuals have a higher metabolic rate than larger (older) ones. The investigations on man clearly show that the difference is due to age and not to size. Age is also probably the determining factor in the metabolic difference found between large and small species, since, in general, species which grow to a large size must be older by the time they have attained that size than are adult individuals of small species.

How far back in the ontogeny can this generalization be carried? At what point in the ontogeny does the metabolic rate attain its highest value? Few researches have been carried out upon these points. The eggs of animals are probably cells of very low metabolic rate. After fertilization, the metabolic rate gradually rises, as development proceeds, up to a certain point and then falls. In the sea-urchin egg (*Arbacia punctulata*),

the oxygen consumption was found to rise continuously up through late cleavage (determinations stopped at this point); and in the eggs of *Fundulus heteroclitus*, up to the time when the embryonic axis is established after which it fell, rising again later apparently as the result of functional activity (unpublished personal observations). Unfortunately in these cases, nothing is known of the metabolic rate of the adult. Hasselbalch ('00) and Bohr and Hasselbalch ('00) have measured the rate of carbon dioxide production and oxygen consumption per kilogram in chick embryos. Both were found to be very high in the earliest stages determined—five or six days,—much higher than those of the adult; the rates then fell rapidly. The carbon dioxide output fell to the ninth day, after which it was about constant, and about equal to that of the adult. The oxygen consumption fell to the eleventh day, then rose again, and on the sixteenth day was again considerably higher than that of the adult. Bohr ('00) compared the carbon-dioxide output of guinea-pig embryos with that of the mother. He did this by cutting open the uterus under anaesthesia, determining the carbon-dioxide output of the mother and foetuses, then clamping the umbilical cords of the foetuses, and again measuring the CO₂ output of the mother. The difference between the two measurements supposedly represents the carbon-dioxide production due to the embryos. Of the six experiments of this kind performed by Bohr, five showed the carbon-dioxide production of the foetuses to be greater than that of the mother. The case where the youngest embryos (5.5 grs. each) were encountered gave a carbon-dioxide output very much higher than that of the mother; the next size (16.5 grs. each) gave less CO₂, but still at a rate considerably higher than that of the mother; while in the other four cases, with large foetuses, the CO₂ production was slightly higher than that of the mother in three, lower in one.¹

¹ I wish to state that the conclusions which I have drawn after careful perusal of the papers of Bohr and Hasselbalch are somewhat different from those stated by the authors. They conclude, curiously enough, that the respiratory rate of the embryos of the hen and guinea pig is no higher than that of the adult, and anyone reading their conclusion alone would certainly be misled. The authors have either overlooked or ignored the fact, which their experimental data clearly show,

These researches indicate that the rate of respiratory exchange is very high at some certain stage in the embryonic development, this stage probably differing in different animals, and falls subsequently. Later it probably rises with increased functional activity. It certainly seems to me that the metabolic rate of embryos, especially vertebrate embryos, cannot validly be compared with that of post-embryonic stages, owing to the enormous differences in functional activity which exist between two such stages. The comparison of the metabolic rate of a chick embryo with an adult hen seems to me a simple absurdity when one reflects upon the difference in muscle tension alone at the two stages. The fact that the total metabolic rate of chick embryos was at no stage found to be less than that of the hen, certainly indicates that if cells of the same degree of functional activity could be compared in embryo and adult, the metabolic rate of the embryonic cells would be vastly the greater. The same criticism applies to cases where the young remain more or less helpless after birth or hatching.

In addition to these researches in which direct measurement of the rate of oxygen consumption or carbon dioxide output of young and old animals have been made, a considerable mass of data is available in which another method was employed. This is the direct susceptibility method extensively used in this laboratory by Child and others; it consists in observing the time of death of organisms in lethal solutions of various substances. We have brought forward a large amount of evidence¹ to indicate that the time of death in such solutions is an index of metabolic rate, individuals of higher metabolic rate dying first. When individuals of different ages are compared by this method, it is invariably found that the time of death is shorter the younger the individual, *always providing that the same degree of functional activity is present in the animals which are being* that the rate of respiratory exchange in the youngest embryos with which they dealt very greatly exceeds that of the adult. It is only at certain later stages that the rates of the two are approximately equal. The authors seem to have an idea that the rate of respiratory exchange ought to be the same throughout development, whereas their own and other data show that it is high in early stages and declines as development proceeds. This decline is probably of the same nature as that which organisms undergo from birth to maturity.

¹ A general résumé of this evidence will be found in Child, '13.

compared. This age difference in susceptibility to toxic solutions has been observed in *Paramecium*, three species of *Hydra*, a number of colonial hydroids and hydromedusæ, several species of planarians, and several small aquatic oligochætes; and the rise in metabolic rate which is a feature of early development has also been demonstrated by this method.

Not only does the rate of respiratory exchange in organisms vary inversely with age but many physiological activities exhibit the same relation. Bert ('70) was among the first to observe this fact. He noted that the rate of respiratory movements is faster in small than in large individuals of the same and related species. Ducceschi ('03) studied the rate of movement of the fins, tail, fin membrane, and operculum of a large number of fish of different sizes; the rate of movement of the maxillipeds, abdominal appendages, and claws of several crustacea; and of the fins and mantle of some cephalopods; and found that it varies inversely as the size of the animal. Mayer ('06) and Child ('18) observed that the rate of contraction of the bells of medusæ is faster the smaller the animal. The number of respiratory movements per minute of *Octopus* is greater the smaller the individual (Polimanti, '13). In four species of sea-cucumber, Crozier ('16) observed that the pulsation of the cloaca is more rapid the smaller the animal. Hecht ('16, '18) noted that small *Ascidia* pass relatively more water through their siphons than large ones; and further that the rate of the heart beat in *Ascidia* varies inversely as the size of the animal. In man, of course, it is well known that the rate of the heart beat is fastest in the fœtus and decreases progressively with age. The rate of growth follows a similar law.

The results of the present experiments are in full accord with those of previous investigators. Small (young) planarians whether asexually or sexually produced, consume more oxygen per unit weight per unit time than large (old) ones.

II. EXPERIMENTS WITH *Planaria dorocephala*.

Planaria dorocephala lives in spring-fed marshes in morainic regions near Chicago. It has never been found sexually mature in nature (although sexual maturity has been experimentally

induced by Dr. Child in this laboratory), but maintains itself by means of asexual fission. In fission, the posterior portion of the body pulls away from the anterior end, regenerates a head and becomes a complete small worm. Such small worms correspond in all tests which have been made upon them, to young worms in species reproducing sexually.

Six experiments were performed in which the rate of oxygen consumption of such small worms, under 10 mm. in length, was compared with that of large worms, over 20 mm. in length. In all cases, both large and small individuals were taken from the same stock and had been kept under the same conditions previous to the experiment. No worms showing signs of recent fission or regeneration were used. The heads were cut off the day before the test was made in order to eliminate movement. Two successive determinations of the oxygen consumption of each lot of worms were usually made, the worms were then weighed, and the oxygen consumption per two hours per 0.5 grams weight then calculated.

The method of determining the rate of oxygen consumption and the method of weighing have been described in previous papers (Hyman, '19a, '19b).

The results of the six experiments are given in Table I. In all cases the smaller worms consume more oxygen per unit weight per unit time than the larger ones. The per cent. difference ranges from 18 to 55, and closer inspection of the table reveals that the amount of difference is correlated with the length of time which has elapsed since the last feeding. In experiment 4, where the small worms respire but 18 per cent. more than the large ones, only two days had elapsed since feeding, while in the other experiments, where three or four days had passed since feeding, the difference is greater. As shown in a preceding paper (Hyman, '19b), feeding greatly increases the rate of oxygen consumption by stimulating the digestive tract. Since the digestive tract is more extensive in large than in small individuals, the difference between such individuals is decreased by feeding. As I did not perform any experiments with worms deprived of food for longer than four days, I do not know whether further starvation would increase the difference between small and large

TABLE I.

COMPARISON OF THE RATE OF OXYGEN CONSUMPTION OF LARGE (OVER 20 MM.) AND SMALL (UNDER 10 MM.) INDIVIDUALS OF *Planaria dorocephala*.

Size of Worm.	C.c. Oxygen Consumed in 2 Hours.	Weight in Grams.	Oxygen Consumed by 0.5 Grs. in 2 Hours.	Per Cent. Difference.
<i>Exp. 1. Worms Collected March 12, Last Fed April 2, Tested April 5.</i>				
<i>Temp. 22.5° C.</i>				
Large.....	0.18 0.20	0.316	0.30	
Small.....	0.24 0.29	0.325	0.40	33
<i>Exp. 2. Worms Collected March 12, Last Fed March 14, Tested March 17.</i>				
<i>Temp. 22° C.</i>				
Large.....	0.12 0.11	0.187	0.30	
Small.....	0.23 0.17	0.255	0.39	30
<i>Exp. 3. Worms Collected Early Winter, Last Fed April 23, Tested April 26.</i>				
<i>Temp. 21° C.</i>				
Large.....	0.09 0.08	0.179	0.23	
Small.....	0.14 0.13	0.216	0.31	48
<i>Exp. 4. Worms Collected March 12, Last Fed April 30, Tested May 2.</i>				
<i>Temp. 21° C.</i>				
Large.....	0.16 0.17	0.160	0.51	
Small.....	0.22 0.24	0.192	0.59	18
<i>Exp. 5. Worms from Mixed Stock, Last Fed May 6, Tested May 10. Temp. 21° C.</i>				
Large.....	0.13 0.12	0.146	0.42	
Small.....	0.17 0.15	0.123	0.65	55
<i>Exp. 6. Worms Collected May 28, Last Fed June 6, Tested June 10.</i>				
<i>Temp. 22° C.</i>				
Large.....	0.33	0.370	0.22	
Small.....	0.33	0.272	0.31	41

worms, but it is scarcely likely since the effect of feeding has almost completely disappeared in four days. In Allen's experiments (Allen, '19) with *Planaria agilis*, worms starved 27 days were used; and in that case the difference between the largest

and smallest worms was about 40 per cent. (Table I., decapitated worms).

III. EXPERIMENTS WITH *Planaria velata*.

Planaria velata lives in the Chicago region in temporary ponds, usually those that are passing into prairie, known to ecologists as "prairie ponds." Like the preceding species it is never found sexually mature, and reproduces exclusively by a peculiar asexual method. After the worms have attained a certain size, pieces drop off from the posterior end, surround themselves with mucus, and pass into an encysted condition. This process continues until the entire worm has formed a series of cysts. Within the cysts, the pieces undergo regeneration into complete worms of very small size, and these emerge from the cysts in about four weeks. This cycle is repeated as long as the ponds contain water; after the water dries up, the cysts remain quiescent until the following spring.

TABLE II.

COMPARISON OF THE RATE OF OXYGEN CONSUMPTION OF LARGE AND SMALL INDIVIDUALS OF *Planaria velata*.

	C.c. Oxygen Consumed in 2 Hours.	Weight in Grams.	Oxygen Consumed by 0.5 Grs. in 2 Hours.
<i>Three Lots of Worms 10-12 mm. Long, before Encystment; Collected March 7, Last Fed March 10, Tested 13. Temp. 22° C.</i>			
Lot C.	0.42 0.52	0.745	0.31
Lot D.	0.58 0.52	0.774	0.35
Lot E.	0.41	0.593	0.34
<i>Three Lots of Worms less than 4 mm. Long, Emerged during May from Cysts Formed by Above Lots of Worms; Fed Several Times after Emergence; Last Feeding June 6, Tested June 9. Temp. 22° C.</i>			
Lot C.	0.16 0.16	0.125	0.64
Lot D.	0.16 0.16	0.127	0.63
Lot E.	0.10 0.09	0.070	0.67

Owing to the nature of the life cycle of this animal, it was not possible to test the large and small worms simultaneously but the large worms were tested before encystment, the small ones after

emerging from the cysts. Three lots of each size were used. The worms were all taken from the same stock. The results are given in Table II. The heads were not removed in these cases, since the worms were also used for another experiment.

The table shows that the small worms consume 100 per cent. more oxygen than the large worms. The greater difference in this case than with the preceding species is probably due to the greater reorganization involved in the production of young worms with this species. As shown in another paper (Hyman, '19c), the process of regeneration of itself brings about a great increase in the rate of oxygen consumption.

IV. EXPERIMENTS WITH *Planaria maculata*.

The so-called species *Planaria maculata* lives in the eastern United States under stones in ponds and in the Chicago region on submersed vegetation. It is highly probable that these two are not the same species as *Planaria maculata* from the Chicago region has never been found sexually mature, while that from Massachusetts is sexually mature and lays capsules all summer long; further the behavior of the two in regeneration is quite distinct. The forms used in this experiment were collected from a pond at Falmouth, Mass., and the experiments were performed at the Marine Biological Laboratory, Woods Hole, Mass., I am indebted to the director, Professor F. R. Lillie, for a research room in this laboratory.

Sexually mature worms, young worms, and egg capsules were collected at Falmouth and brought to Woods Hole. The rate of oxygen consumption of the mature worms was tested with one exception soon after they were collected. The young worms collected, together with those which subsequently emerged from the capsules, were kept for some time and fed at short intervals on liver (mostly fish liver) until a considerable number of them were at hand, whereupon their rate of oxygen consumption was determined. In all cases the heads were removed at least several hours before the test.

The results are recorded in Table III. The young worms consume about 50 per cent. more oxygen than the sexually mature individuals.

TABLE III.

COMPARISON OF THE RATE OF OXYGEN CONSUMPTION OF SEXUALLY MATURE INDIVIDUALS (15 MM. OR LONGER) WITH THE SEXUALLY PRODUCED YOUNG (5 MM. OR LESS) OF *Planaria maculata*.

No. of Lot.	C.c. Oxygen Consumed in Test.	Weight in Grams.	Oxygen Consumed. by 0.5 Grams in 2 Hours.
<i>Four Lots of Sexually Mature Individuals, Collected July 10; First Three Lots Tested July 11; Fourth Lot Kept Until August 11, with Frequent Feedings, Last Feeding August 8, Tested August 11. Temp. 20° C.</i>			
Lot 1.....	0.30 in 2 0.31 hours	0.700	0.21
Lot 2.....	0.32	0.677	0.23
Lot 3.....	0.32 0.30 0.26	0.680	0.20
Lot 4.....	0.09 0.07	0.152	0.17
<i>Three Lots of Sexually Produced Young; Young and Capsules Collected July 10 Fed at Frequent Intervals; Last Feeding of Lots 1 and 2, July 28, Tested July 31; Lot 3, Last Feeding August 8, Tested August 11. Temp. 20° C.</i>			
Lot 1.....	0.14 in 3 0.12 hours	0.101	0.32
Lot 2.....	0.11	0.091	0.32
Lot 3.....	0.11 0.16 0.18	0.117	0.28

V. CONCLUSIONS.

These experiments show that small or young planarians consume oxygen at a faster rate per unit weight than larger or older ones. As already stated, the carbon dioxide production is also inversely proportional to the size of the worms. It is true that experiments of this kind do not and cannot prove that the protoplasm of young animals actually has an intrinsically higher metabolic rate than that of older ones, for the reason that it is impossible to discover what part of the weight of an animal is active protoplasm and what part inert material. Nevertheless there cannot be any reasonable doubt that the metabolic rate is inversely proportional to age. It would be difficult to suggest any other explanation for many of the facts cited in this paper, namely, for the faster rate of respiration, faster heart beat, and more rapid rate of other physiological activities of young as

compared with older organisms. The fact further that the susceptibility of young animals to a number of toxic substances is greater than that of old could scarcely be supposed to be due to a greater percentage of inert materials in the older individuals,

An interesting point brought out in these experiments is that the difference between the asexual and the sexual young and their respective adults is of about the same magnitude, when considered the same length of time after feeding. Worms produced by fission are therefore as truly "young" as those which develop from the egg.

In previous papers of this series (Hyman, '19b, '19c), it was shown that planarians which have been starved seven or eight weeks and pieces of planaria which have undergone regeneration have a much higher rate of oxygen consumption than ordinary fed worms, all tests being made, of course, a few days after feeding. Starved, regenerated and young worms therefore have this physiological characteristic in common: their metabolic rate is higher than that of large fed worms. That of starved ones is highest, regenerated ones next, and young, when produced from the egg or simple fission, least. It therefore appears that the metabolic rate of reduced forms depends primarily upon the amount of reorganization involved in their production, and is proportional to the degree of reorganization which has taken place. As a further illustration of this may be cited the much higher metabolic rate of the asexual young of *Planaria velata* than those of *P. dorocephala*, presumably because much more extensive changes are involved in giving rise to the former. The evidence presented in these papers clearly supports the view which has been long maintained by Child—that such reorganizations due to whatever cause are rejuvenating transformations, restoring the organism to a physiological condition resembling that of the young.

V. SUMMARY.

1. The young of *Planaria dorocephala* produced by simple fission were found to consume 15 to 55 per cent. more oxygen than large worms, the difference depending upon the length of time which had elapsed between the last feeding and the time of testing.

2. The young of *Planaria velata*, produced by an asexual process involving regeneration and a high degree of reorganization of the body, consume about 100 per cent. more oxygen than the worms from which they come. The greater difference in this species is undoubtedly associated with the method by which the young are produced.

3. The sexually produced young of *Planaria maculata* consume about 50 per cent. more oxygen than the sexually mature worms. There is thus no significant difference between sexual and asexual young, when the latter arise by ordinary simple fission.

4. This result, that young worms have a higher metabolic rate than old ones, is in accord with a considerable body of literature on other forms leading to the same conclusion; and confirms the work previously done in this laboratory upon this same point by other methods.

Allen, G. D.

- '19 Quantitative Studies on the Rate of Respiratory Metabolism in Planaria. II. The Rate of Oxygen Consumption during Starvation, Feeding, Growth, and Regeneration in Relation to the Method of Susceptibility to Potassium Cyanide as a Measure of Rate of Metabolism. *Am. Journ. Physiol.*, Vol. 49, pp. 420-474.

Bert, P.

- '70 Leçons sur la physiologie comparée de la respiration. Paris.

Benedict, F. G., and Talbot, F. B.

- '14 The Gaseous Metabolism of Infants. *Carn. Instit. Wash. Publ. no. 201*, 168 pp.

- '15 The Physiology of the New-born Infant. *Ibid.*, no. 233, 126 pp.

Bohr, Chr.

- '00 Der respiratorische Stoffwechsel des Säugethierembryo. *Skand. Arch. f. Physiol.*, Vol. 10, pp. 412-433.

Bohr, Chr., and Hasselbalch, K. A.

- '00 Ueber die Kohlensäureproduktion des Hühnerembryos. *Skand. Arch. f. Physiol.*, Vol. 10, pp. 149-173.

Bounhiol, J.

- '02 Recherches biologiques expérimentales sur la respiration des Annélides polychètes. *Ann. d. Sci. Nat. Zool.*, Vol. 16, pp. 1-128.

Child, C. M.

- '13 Studies on the Dynamics of Morphogenesis and Inheritance in Experimental Reproduction. V. The Relation Between Resistance to Depressing Agents and Rate of Metabolism in *Planaria dorotocephala* and its Value as a Method of Investigation. *Journ. Exp. Zool.*, Vol. 14, pp. 153-206.

- '18 Physiological Senescence in *Hydromedusa*. *Biol. Bull.*, Vol. 34, pp. 49-63.

- '19 A Comparative Study of Carbon Dioxide Production during Starvation in Planaria. *Am. Journ. Physiol.*, Vol. 48, pp. 231-257.

Crozier, W. J.

- '16 The Rhythmic Pulsation of the Cloaca of Holothurians. Journ. Exp. Zööl., Vol. 20, pp. 297-356.

Du Bois, E. F.

- '16 Clinical Calorimetry. XII. The Metabolism of Boys' Twelve and Thirteen Years Old compared with the Metabolism at Other Ages. Archives Int. Med., Vol. 17, pp. 887-901.

Ducceschi, V.

- '03 Una legge del movimento animale. Zeitsch. f. allg. Physiol., Vol. 2, pp. 482-501.

Gephart, F. C., and Du Bois, E. F.

- '16 Clinical Calorimetry. XIII. The Basal Metabolism of Normal Adults with Special Reference to Surface Area. Archives Int. Med., Vol. 17, pp. 902-914.

Hasselbalch K. A.

- '00 Ueber den respiratorischen Stoffwechsel des Hühnerembryos. Skand. Arch. Physiol., Vol. 10, pp. 353-402.

Hecht, S.

- '16 The Water Current Produced by *Ascidia atra* Lesueur. Journ. Exp. Zööl., Vol. 20, pp. 429-434.
'18 The Physiology of *Ascidia atra* Lesueur. III. The Blood System. Am. Journ. Physiol., Vol. 45, pp. 157-187.

Hill, A. V. and A. M.

- '13 Calorimetrical Experiments on Warm-blooded Animals. Journ. of Physiol., Vol. 46, pp. 81-103.

Hyman, L. H.

- '19a On the Action of Certain Substances on Oxygen Consumption. II. Action of Potassium Cyanide on Planaria. Am. Journ. Physiol., Vol. 48, pp. 340-371.
'19b Physiological Studies on Planaria. I. Oxygen Consumption in Relation to Feeding and Starvation. Am. Journ. Physiol., Vol. 49, pp. 377-402.
'19c Idem. II. Oxygen Consumption in Relation to Regeneration. Ibid., Vol. 50, pp. 67-81.

Jolyet, F., and Regnard, P.

- '77 Recherches physiologiques sur la respiration des animaux aquatiques. Arch. d. Physiol., 2d Ser., Vol. 4, pp. 44, 584.

Magnus-Levy, A., and Falk, E.

- '99 Der Lungengaswechsel des Menschen in der verschiedenen Altersstufen. Arch. f. Physiol., Suppl. Bund., pp. 314-381.

Mayer, A. G.

- '06 Rhythmical Pulsations in *Scyphomedusa*. Carn. Instit. Wash. Publ. no. 47, 62 pp.

Montuori, A.

- '13 Les processus oxydatifs chez les animaux marins en rapport avec la loi de superficie. Arch. Ital. d. Biol., Vol. 59, pp. 213-234.

Morgulis, S.

- '15 The Body Surface of Flounders and its Relation to the Gaseous Metabolism. Am. Journ. Physiol., Vol. 36, pp. 207-216.

Nicolas, M. G.

- '18 Contribution à l'étude des variations de la respiration des végétaux avec l'âge. *Rev. gen. d. Bot.*, Vol. 30, pp. 209-225.

Polimanti, O.

- '13 Sui rapporti fra peso del corpo e ritmo respiratorio in *Octopus vulgaris* Lam. *Zeitschr. f. allg. Physiol.*, Vol. 15, pp. 449-455.

Slowtsoff, B.

- '03 Ueber die Beziehungen zwischen Körpergrösse und Stoffverbrauch der Hunde bei Ruhe und Arbeit. *Arch. f. d. ges. Physiol.*, Vol. 95, pp. 158-191.

Tashiro, S., and Adams, H. S.

- '14 Comparison of the Carbon Dioxide Output of Nerve Fibers and Ganglia in *Limulus*. *Journ. Biol. Chem.*, Vol. 18, pp. 329-334.

Vernon, H. M.

- '95 The Respiratory Exchange of the Lower Invertebrates. *Journ. of Physiol.*, Vol. 19, pp. 18-70.

ON THE ACTION OF CERTAIN SUBSTANCES ON OXYGEN CONSUMPTION.

III. ACTION OF POTASSIUM CYANIDE ON SOME CŒLEENTERATES AND ANNELIDS.

L. H. HYMAN,

HULL ZOÖLOGICAL LABORATORY, UNIVERSITY OF CHICAGO.

In this paper are presented some further data concerning the effect of potassium cyanide on the rate of oxygen consumption of animals. In view of the importance of this substance as a reagent in physiological experiments, and since tests of this kind have been made upon comparatively few animals, it has seemed worth while to obtain data upon some representatives of groups in which the action of cyanide has not yet been determined. No experiments have, to my knowledge, been performed upon annelids, and only one cœlenterate has been tested, namely *Gonionemus*, in which form Loeb and Wasteneys ('13) showed that potassium cyanide decreases the oxygen consumption to a considerably greater extent than does ethyl urethane, although both produce the same degree of anæsthesia.

The literature dealing with the chemical, physiological and pharmacological properties of the cyanides and related substances has been extensively reviewed in the second paper of this series (Hyman, '19), and will therefore not be restated here. Briefly, it may be said that this group of substances has been shown to depress many physiological processes; and to decrease the rate of oxygen consumption or carbon dioxide output or both of the following living materials: yeast, a mould *Aspergillus*, a sponge *Suberites*, *Gonionemus*, *Planaria*, a beetle *Passalus cornutus*, embryos of *Fundulus*, several mammals, sea-urchin eggs, red blood corpuscles of geese, isolated mammalian kidneys, the frog's heart, minced beef liver, minced horse, beef, and pigeon muscle, and minced horse brain. In most of these cases it was shown that the effect was reversible. Since that review was

written, two additional papers have come to my attention, those of Allen ('19) and of Evans ('19). Allen's results and conclusions regarding the effect of potassium cyanide on the rate of oxygen consumption of *Planaria* are about the same as my own, except for one or two points; thus Allen states that the oxygen consumption of *Planaria* cannot be reduced to less than 20 per cent. of the normal by cyanide, while I found in many cases, reduction to 10 to 15 per cent. of the normal. Evans's interesting paper contains some further references to the literature, confirms some previous work, and presents a number of new facts. Thus Evans finds that cyanide decreases markedly and reversibly the oxygen consumption of the cat, confirms the older statements that oxygen is just as readily dissociable from the blood in cyanide poisoning as from normal blood; and shows a striking similarity between the behavior of smooth, skeletal, and heart muscle, and nerve and nervous centers in the presence of cyanide and in lack of oxygen. Evans concludes that the cyanides "appear to exert all of their physiological effects by reason of the anoxæmia which they produce," and favors the view that they act by uniting chemically with some tissue constituent.¹

¹ Comment would seem to be required on Evans's remark (p. 23) that the observations of Lund and Herwerden seem to be incompatible with Child's conception of metabolic gradients. Lund's results have been considered elsewhere (Child, '19). Herwerden ('18) says that the buds and young of *Hydra* are not more susceptible to cyanide than the parents. Perusal of her paper shows, however, that she has compared only three or four pairs of individuals and has observed the *disintegration of the tentacles only*, having removed the cyanide when this occurred. Now, we have clearly shown (Child and Hyman, '19) that the buds of *Hydra* are not comparable with the parents *until they have the same degree of muscular activity*, and further that the susceptibility of the tentacles is also *dependent upon their degree of activity*. Miss Herwerden is mistaken in her conclusion. We have compared dozens of young and adult *Hydras*, belonging to three species, and have always found that the young are more susceptible to cyanide *when comparable degrees of muscular activity* exist. It is true that the difference is *least* in the case of the tentacles but even there the tentacles of the young are slightly more susceptible. Neither are Miss Herwerden's statements about the susceptibility of the Daphnid heart entirely correct. I have found that the smallest individuals in a *Daphnia* culture are the most susceptible of all (both as to heart and entire animal); that the susceptibility decreases with increasing size up through the size when the females are producing their first parthenogenetic broods; but from that time on, the relation is reversed, so that the largest and oldest individuals in the culture, which have produced many parthenogenetic broods, are more susceptible than

In the experiments to be reported in this paper it was not my purpose to make a complete study of the action of cyanide on oxygen consumption in these animals, since I had already done that in the case of a sponge and *Planaria*, but merely to show that cyanide in non-lethal doses reversibly decreases oxygen intake. The experiments on the marine forms were performed at the Marine Biological Laboratory, Woods Hole, Mass.; those on the fresh-water forms at the University of Chicago. I am greatly indebted to Professor F. R. Lillie for working space at the first-named institution.

The methods employed were identical with those used in previous experiments on this subject and will be found described in full in a former paper (Hyman, '19). Briefly, the animals to be tested were placed in Erlenmeyer flasks or wide-mouthed bottles of 500 to 600 c.c. capacity, these were filled air-tight with water, and the animals allowed to respire in them for a convenient length of time. A sample was then withdrawn, and this and a sample of the original water used were analyzed for oxygen content by Winkler's method, the difference between the two samples representing the oxygen consumed by the animals. Two separate determinations of the oxygen consumption in normal water were made, potassium cyanide was then added to the water and two more determinations in the presence of cyanide carried out. The animals were then washed in several changes of water, and their oxygen consumption in normal water tested again a day or two later. Throughout each experiment, the receptacles containing the animals were immersed in a water-bath, the temperature of which was kept constant to 0.5 degrees.

The possibility of iodine absorption by substances emanating from the organisms was again considered, and was tested in the case of two or three of the animals employed. Standard iodine solution was added to water which had been standing on the animals for at least one hour, and the iodine then titrated back

half-grown individuals. It would thus appear that in these animals parthenogenetic reproduction is accompanied by some degree of rejuvenescence. Green's recent work on *Simocephalus* (BIOL. BULL., Aug., 1919) supports this suggestion, since he found that these animals reproduce sexually early in ontogeny, and subsequently the same individuals begin parthenogenetic reproduction, a reversal of the order usually observed in parthenogenetic forms.

with thiosulphate. No loss of iodine was found, and indeed, this possible source of error in the Winkler method would seem to have been greatly overrated (*cf.* Hale and Melia, '13).

EXPERIMENTS ON PENNARIA.

Pennaria tiarella, a common colonial hydroid of the Atlantic coast, was selected as a representative of this group of cœlenterates. Large quantities of freshly collected material were available. Clean branches, free from visible plant growth, were cut off, *Caprella* and other small animals which commonly live on such colonies carefully removed, and the branches were then placed in wide-mouthed bottles, and treated as described above. The animals were always used within a few hours after they were collected.

TABLE I.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF
Pennaria tiarella.

No. of Experiment	1.	2.	3.	4.	5.	6.
Temp. and Date	Aug. 18, 19 ^o C.			Aug. 23, 22 ^o C.		
<i>Oxygen Consumed, Cubic Centimeters per Hour.</i>						
First hour normal.	0.55	0.56	0.43	0.94	0.78	1.21
Second hour normal.	0.62	0.56	0.43	0.95	0.74	1.20
	1/25000 Mol. KNC.			1/10000 Mol. KNC.		
First hour KNC.	0.38	0.38	0.27	0.52	0.45	0.60
Second hour KNC.	0.32	0.35	0.25	0.36	0.34	0.54
Per cent. decrease.	41	35	39	54	49	53

Six different lots of *Pennaria* were tested in this manner, as to their normal rate of oxygen consumption and the rate in 1/10000 and 1/25000 mol. potassium cyanide. The results are given in Table I. In all cases, the rate of oxygen consumption is reduced in the presence of cyanide. Unfortunately in the case of this animal it could not be determined whether the effect was reversible, since *Pennaria* deteriorates rapidly in the laboratory; within twenty-four hours, the majority of the hydranths have either fallen off or have lost their normal appearance.

Since *Pennaria* exhibits very few movements, it is not possible

that differences in muscular activity could account for the decreased oxygen intake in the presence of cyanide. As far as could be determined, the hydranths were as fully expanded in the presence of cyanide as in normal sea-water.

EXPERIMENTS ON METRIDIDIUM.

Metridium marginatum was selected as a representative of the Anthozoa. The individuals used had probably been kept in float cars for some time. Medium-sized individuals were placed in wide-mouthed bottles of about 600 c.c. capacity, one in each, and left there undisturbed until they had attached themselves and expanded fully. It was found that the animals soon became accustomed to such an environment and could be handled and would submit to change of water without contracting. Since the oxygen consumption of sea-anemones in all probability varies with the degree of expansion, note was always made of the degree of expansion during the experiments. In experiments 1 and 3, the animals were slightly contracted during the exposure to cyanide but in the other six experiments they were fully expanded throughout. Differences in state of muscular activity are therefore not responsible for the observed results.

The eight experiments which were performed upon *Metridium* are reported in Table II. A marked and reversible decrease in oxygen consumption in the presence of cyanide was found in all cases.

TABLE II.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF
Metridium marginatum.

No. of Experiment,	1.	2.	3.	4.	5.	6.	7.	8.
Temp. and Date	Aug. 12, 21° C.			Aug. 10, 21.5° C.		Aug. 13, 20° C.		
<i>Oxygen Consumed, Cubic Centimeters per Hour.</i>								
First hour normal.	1.33	1.19	1.13	1.66		1.06	1.01	0.97
Second hour normal.	1.09	1.05	1.03	1.94	1.67	1.12	1.19	1.31
	1/5000 Mol. KNC.			1/10000 Mol.		1/25000 Mol.		
First hour KNC.	0.48	0.56	0.56	1.24	1.25	0.81	0.70	0.66
Second hour KNC.	0.49	0.59	0.42	0.89	0.70	0.70	0.76	0.59
Per cent. decrease.	60	49	55	41	58	32	34	46
Recovery.	1.09	1.10	1.14	1.68		1.08	1.18	1.20

EXPERIMENTS ON POLYCHÆTES.

Experiments were performed on three common polychæte annelids of the Atlantic coast—*Nereis virens*, *Arenicola cristata*, and *Chætopterus pergamentaceus*. Of the three species, *Chætopterus* was found to be the most favorable for this kind of experimentation, as it is not a very active animal. No difference in amount of activity in normal and cyanide-containing sea-water was observed. *Arenicola* was fairly favorable for the purpose as it also is relatively inactive, and here again there was no significant difference in activity throughout the experiments; indeed, it seemed to me that the respiratory movements were more pronounced in the presence of cyanide than in normal sea-water. On the other hand, the results with *Nereis* were not very satisfactory owing to the restlessness of the animals. In experiments 1-3, Table V, the animals were considerably more active in normal sea-water than they were after cyanide had been added, and hence the decrease observed in cyanide was due in part to diminished motor activity. In experiments 4-6, Table V., however, differences in motor activity were not present and have not contributed to the result.

A further difficulty was encountered in the case of *Nereis*. It was observed that both the degree of activity and the rate of oxygen consumption of *Nereis* diminished when it was kept in the laboratory. Experiments 1-3 were performed on freshly collected individuals and it will be noted that the oxygen consumption of these individuals is much higher than in the other cases, and further that the recovery figures, obtained upon them twenty-four hours later, are considerably lower than the original figures. This is not due to the exposure to cyanide since individuals from the same collection which had not been subjected to cyanide showed the same decrease in oxygen intake after a day in the laboratory. It is probable that this decrease is due in large part to starvation. I have observed it in the case of a number of other animals also, and my experience indicates that animals which have been kept for a few days after removal from their natural environment are preferable to freshly collected material for experiments of this kind. This difficulty was not encountered in the case of *Chætopterus* and *Arenicola* since it happened that

the animals had been on hand in the collecting department for some time before I obtained them.

Since the first experiments showed that some little time was required for the penetration of even relatively concentrated

TABLE III.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF
Chatopterus pergamentaceus.

No. of Experiment	1.	2.	3.	4.	5.	6.	7.	8.	6.
Temp. and Date	Aug 3, 20.5° C.			Aug. 3, 21° C.			Aug. 5, 21° C.		
<i>Oxygen Consumed, Cubic Centimeters per Hour.</i>									
First hour normal.....	0.48	0.40	0.56	0.44	0.36	0.45	0.47	0.35	0.44
Second hour normal.....	0.33	0.40	0.48	0.40	0.36	0.44	0.49	0.32	0.43
	<i>1/2000 Mol. KNC (Two Hrs. Before Test).</i>			<i>1/5000 Mol. KNC (Two Hrs. Be- fore).</i>			<i>1/10000 Mol. KNC (One Hr. Be- fore).</i>		
First hour KNC.....	0.20	0.13	0.23	0.23	0.16	0.27	0.41	0.24	0.34
Second hour KNC.....	0.14	0.16	0.21	0.14	0.16	0.22	0.41	0.22	0.33
Per cent. decrease.....	59	64	58	56	56	45	15	32	23
Recovery.....	0.34	0.48	0.50	0.48	0.34	0.43			

TABLE IV.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF
Arenicola cristata.

No. of Experiment	1.	2.	3.	4.	5.	6.
Temp. and Date	Aug. 2, 20.5° C.			Aug. 1, 21° C.		
<i>Oxygen Consumed, Cubic Centimeters per Hour.</i>						
First hour normal.....	0.42	0.39	0.67	0.42	0.48	0.43
Second hour normal.....	0.34	0.38	0.63	0.44	0.47	0.56
	<i>1/1000 Mol. KNC (Two Hrs. Before).</i>			<i>1/2000 Mol. KNC (Two Hrs. Before).</i>		
First hour KNC.....	0.27	0.30	0.46		0.22	0.31
Second hour KNC.....	0.20	0.26	0.46	0.27		0.30
Per cent. decrease.....	39	28	30	35	54	39
Recovery.....	0.49	0.31	0.71	0.42	0.48	0.43

solutions of cyanide into these animals, presumably because of their thick body walls, the practice of leaving them for an hour or two in the cyanide solution before carrying out the test was adopted.

The results with *Chaetopterus* are presented in Table III., with *Arenicola* in Table IV, and with *Nereis* in Table V. One or two individuals, depending on size, were placed in each flask. For reasons already given, experiments 1-3, Table V., on *Nereis* were not very satisfactory but the other experiments on these polychaetes leave no doubt that cyanide brings about a reversible decrease in oxygen consumption.

TABLE V.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF
Nereis virens.

No. of Experiment	1.	2.	3.	4.	5.	6.
Temp. and Date	Aug. 5, 22° C.			Aug. 6, 21° C.		
<i>Oxygen Consumed, Cubic Centimeters per Hour.</i>						
First hour normal	0.80	0.65	0.65	0.30	0.36	0.29
Second hour normal	0.78	0.55	0.64	0.23	0.23	
	1/5000 Mol. KNC (Two Hrs. Before Test),			1/1000 Mol. KNC (One-half Hour Before).		
First hour KNC	0.14	0.11	0.11	0.05	0.05	0.08
Per cent decrease	83	82	83	83	84	73
Recovery	0.32 ¹		0.48 ¹	0.34		0.29

EXPERIMENTS ON LEECHES.

The leeches used in these experiments were *Hamopsis marmoratis*, and *Herpobdella punctata*, chiefly the latter. They were obtained from ditches near Wolf Lake, Indiana, and subjected to experiment shortly after they were brought into the laboratory. Two or three individuals were placed in each flask. As in the case of the polychaetes, it was impossible to eliminate movement, but the degree of activity was about the same in both normal and cyanide-containing water. The results are given in Table VI. Experiments 4 and 7 were performed upon *Hamopsis*, the others on *Herpobdella*.

¹ Failure to recover original rate of oxygen consumption not due to cyanide, see text.

TABLE VI.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF LEECHES
(*Hamopsis marmoratis*, EXPS. 4 AND 7, AND *Herpobdella punctata*,
REMAINING EXPS.).

No. of Experiment	1.	2.	3.	4.	5.	6.	7.	8.	9.	10.
Temp. and Date	April 3, 21° C.			Mar. 22, 21° C.			April 1, 21° C.			
<i>Oxygen Consumed, Cubic Centimeters per Hour.</i>										
First hour normal	0.15	0.22	0.19	0.51	0.31	0.28	0.51	0.21	0.24	0.20
Second hour normal	0.16	0.24		0.51	0.30	0.25	0.48			
	1/5000 Mol. KNC.			1/10000 Mol. KNC.			1/25000 Mol. KNC.			
First hour KNC	0.01	0.07	0.04	0.17	0.17	0.15	0.29	0.08	0.17	0.14
Second hour KNC	0.04	0.07	0.06	0.13	0.14	0.11	0.26	0.09	0.14	0.13
Per cent. decrease	84	70	74	73	51	51	45	60	36	33
Recovery	0.17	0.28	0.23		0.26	0.33		0.16	0.23	0.19

EXPERIMENTS ON OLIGOCHÆTES.

The forms used in these experiments were an aquatic earth-worm, *Helodrilus tetædra*, one of the megadrilous oligochætes, and *Lumbriculus inconstans*, one of the microdrilous oligochætes. Both of these annelids live in temporary pools in the woods in the dune region of Indiana; the former occurs in the larger pools and also in permanent ponds, while the latter species has never been found in permanent bodies of water. They were collected near Clarke, Indiana. Both species are highly thigmotactic, being found in their natural habitat entwined among the branches of the moss which commonly grows in these pools or lying between layers of dead leaves, their posterior ends usually protruded for respiratory purposes. As the animals are very restless when their bodies are not in contact with objects, an attempt was made to quiet them by placing a small amount of thoroughly washed cotton in the experimental flasks with the worms for some hours preceding the tests. This procedure was entirely successful with *Helodrilus*; the worms entwined themselves among the cotton fibers and remained perfectly quiet throughout the experiments. It was possible to quiet the majority of the individuals of *Lumbriculus* in this way also, but a few individuals would always continue to crawl about. No difference, however,

was observed in the degree of activity in normal and in cyanide-containing water.

The results with *Helodrilus* are presented in Table VII. and with *Lumbriculus* in Table VIII. As these worms are rather

TABLE VII.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF
Helodrilus tetædra.

No. of Experiment	1.	2.	3.	4.	5.	6.
Temp. and Date	June 3, 23° C.			June 2, 23° C.		
<i>Oxygen Consumed, Cubic Centimeters in Two Hours.</i>						
First hour normal	0.13	0.16	0.16	0.17	0.18	0.19
Second hour normal	0.13	0.19	0.15	0.16	0.22	0.22
	1/5000 Mol. KNC.			1/10000 Mol. KNC.		
First hour KNC	0.05	0.08	0.09	0.13	0.16	0.11
Second hour KNC	0.06	0.06	0.07	0.09	0.12	0.08
Per cent. decrease	59	60	49	30	30	54
Recovery	0.14	0.16		0.14	0.19	0.16

TABLE VIII.

ACTION OF POTASSIUM CYANIDE ON THE OXYGEN CONSUMPTION OF
Lumbriculus inconstans.

No. of Experiment	1.	2.	3.	4.	5.	6.
Temp. and Date	Oct. 17, 21° C.			Oct. 15, 21° C.		
<i>Oxygen Consumed, Cubic Centimeters per Hour.</i>						
First hour normal	0.44	0.47	0.37	0.58	0.58	0.46
Second hour normal	0.45	0.41	0.29(?)	0.57	0.61	0.49
	1/5000 Mol. KNC.			1/10000 Mol. KNC.		
First hour KNC	0.20	0.24	0.18	0.30	0.28	0.21
Second hour KNC	0.14	0.16	0.14	0.28	0.26	0.19
Per cent. decrease	62	55	52	50	55	56
Recovery	0.38	0.42	0.42			

small, a large number of individuals was used in each experiment. The results are the same as in the case of the other animals tested, a reversible decrease in oxygen intake when cyanide is present.

SUMMARY AND CONCLUSIONS.

1. The normal oxygen consumption and the oxygen consumption in the presence of various concentrations of potassium cyanide was tested in the case of *Pennaria*, *Metridium*, *Nereis*, *Chaetopterus*, *Arenicola*, two species of leeches, and two species of aquatic oligochaetes. In all cases, numbering about seventy experiments (not all of which are reported), the oxygen consumption was markedly decreased in the presence of cyanide.

2. This decrease was not due to differences in muscular activity, since in some of the experiments it was possible to keep the animals entirely quiet throughout, while in the others, with three exceptions noted in the text, the animals were equally active in both the normal and the cyanide-containing water. None of the concentrations of cyanide used caused any visible anæsthesia, within the short time periods during which the animals were exposed to them.

3. The decrease was reversible, the oxygen consumption returning to approximately the original value when the cyanide was washed out of the animals. In no case were the animals injured in any way. Most of them were kept for a considerable length of time after the experiments were completed and were entirely normal in behavior and appearance.

4. As was found to be the case in previous experiments with cyanide, the percentage of decrease is absolutely greater, the more concentrated the cyanide solution, but the more dilute solutions are relatively more effective.

5. These results are in accord with previous experiments on the action of cyanides and justify the use of the cyanides as depressing agents.

LITERATURE.

Allen, G. D.

- '19 Quantitative Studies on the Rate of Respiratory Metabolism in *Planaria*.
I. The Effect of Potassium Cyanide on the Rate of Oxygen Consumption.
Amer. Jour. Physiol., Vol. 48, pp. 93-121.

Child, C. M.

- '19 A Comparative Study of Carbon Dioxide Production during Starvation in *Planaria*. Amer. Jour. Physiol., Vol. 48, pp. 231-257.

Child, C. M., and Hyman, L. H.

- '19 The Axial Gradients in Hydrozoa. I. Hydra. BIOL. BULL., Vol. 36, pp. 183-223.

Evans, C. Lovatt.

'19 Observations on Cyanide Anoxæmia. *Jour. of Physiol.*, Vol. 53, pp. 17-42.

Hale, F. E., and Melia T. W.

'13 Winkler's Method for the Determination of Oxygen in Water; the Effect of Nitrite and its Prevention. *Jour. Indust. and Engineer. Chem.*, Vol. 5, pp. 976-980.

Herwerden, M. A. van.

'18 La résistance de cellules de divers âges a l'empoisonnement par le cyanure de potassium. *Arch. neerl. de Physiol.*, Vol. 2, pp. 715-720.

Hyman, L. H.

'19 On the Action of Certain Substances on Oxygen Consumption. II. Action of Potassium Cyanide on *Planaria*. *Am. Jour. Physiol.*, Vol. 48, pp. 340-371.

Loeb, J., and Wasteneys, H.

'13 Narkose und Sauerstoffverbrauch. *Biochem. Zeitschr.*, Vol. 56, pp. 295-306.

MBL/WHOI LIBRARY



WH 17K9 0

